

Anti-MTA1 Antibody Picoband®

Catalog Number: PA1483

About MTA1

Metastasis-associated protein MTA1 is a protein that in humans is encoded by the MTA1 gene. This gene encodes a protein that was identified in a screen for genes expressed in metastatic cells, specifically, mammary adenocarcinoma cell lines. Expression of this gene has been correlated with the metastatic potential of at least two types of carcinomas although it is also expressed in many normal tissues. By fluorescence in situ hybridization, mapped the MTA1 gene to chromosome 14q32.3. MTA1 is a component of the chromatin remodeling complex that influences gene transcription by modulating target gene chromatin. MTA1 is widely upregulated in many carcinomas.

Overview


Product Name	Anti-MTA1 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-MTA1 Antibody catalog # PA1483. Tested in IHC, IHC-F, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, IHC-F, WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q13330

Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human MTA1, identical to the related mouse and rat sequences.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and IHC(F).
Cross Reactivity	No cross-reactivity with other proteins

Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Immunohistochemistry (Frozen Section), 0.5-1ug/ml, Mouse, Rat, Human Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, Mouse, Rat Western blot, 0.1-0.5ug/ml, Human, Rat, Mouse


Anti-MTA1 Antibody Picoband® (PA1483) Images



Western blot analysis of MTA1 using anti-MTA1 antibody (PA1483). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human MCF-7 whole cell lysates, Lane 2: human HEK293 whole cell lysates, Lane 3: human HELA whole cell lysates, Lane 4: human PC-3 whole cell lysates, Lane 5: human Raji whole cell lysates, Lane 6: human K562 whole cell lysates, Lane 7: rat brain tissue lysates, Lane 8: mouse brain tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MTA1 antigen affinity purified polyclonal antibody (Catalog # PA1483) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for MTA1 at approximately 80KD. The expected band size for MTA1 is at 80KD.






The expression of MTA1 is up-regulated in RCCs. a Immunohistochemistry was used to characterize the expression of MTA1 in RCC and adjacent tissues. MTA1 was highly expressed in ccRCCs, compared to very weak staining in adjacent tissue. b Image Pro Plus was used for the statistical analysis of the positive signal (IOD) of MTA1 expression in ccRCCs and the adjacent tissue. The expression of MTA1 was significantly up-regulated in ccRCCs. ** p




MTA1 promotes A498 cell migration. A498 cells were transfected with pcDNA3.1-Flag, Flag-MTA1, siNC, and si-MTA1. After 24 h, cells in the four conditions were subjected to the wound healing assay. Macrographs were taken under $\times 100$ magnification Index in PubMed under a CC BY license. PMID: 33059651

MTA1 enhances the invasion of A498 cells. a A498 cells were transfected with pcDNA3.1-Flag, Flag-MTA1, siNC, and si-MTA1. After 24 h, the transwell cell invasion assay using A498 cells was performed, and macrographs were taken under $\times 40$ magnification. Scale bar: 50 μ m. b Image J was used to count the transmigrated cells and Student's t test to




analyze differences. ** p


MTA1 regulated the expression of metastasis-related factors via the NFkappaB pathway. A498 cells were transfected with pcDNA3.1-Flag, Flag-MTA1, siNC, and si-MTA1. a Protein levels of MTA1; cells were collected for WB with MTA1 and GAPDH antibodies. The bands of over-expressed MTA1 were from Additional file : Fig. 2A and down-expressed MTA1 were from Additional file : Fig. 2B with arrow. b Cells were lysed and used in qRT-PCR assays to measure the E-cadherin mRNA expression. Statistical analysis was performed using Student's t test. b , c A498 cells were transfected with pcDNA3.1-Flag and Flag-MTA1. After 36 h, the Flag-MTA1 group was treated with 10 nM PDTC for 12 h. Then cells were harvested for qRT-PCR evaluation to measure the mRNA expression of MMP2 (b) and MMP9 (c). Statistical analysis was performed using ANOVA. d A498 cells were transfected with pcDNA3.1-Flag, Flag-MTA1, siNC, and si-MTA1. After 48 h, cells were subjected to western blotting analysis using antibodies targeting p-p65, p65 and ACTB. The bands of p-p65, p65 and ACTB were from Additional file : Fig. 3. e, f Image J was used to calculate the gray scanned bands of p-p65 (b) and p65 (c). * p




MTA1 regulated the migration and invasion of RCC cells via NF-kappaB. A498 cells were transfected with pcDNA3.1-Flag, Flag-MTA1, and Flag-MTA1 + PDTC (10 nM). a In the wound healing assay, MTA1 improved the migration of RCC cells, while the addition of the inhibitor PDTC blocked the effect of MTA1 on migration. Scale bar: 100x. b In the transwell assay, MTA1 markedly induced invasion of A498 cells, but compared to the MTA1 group, the invasion of MTA1 + PDTC A498 cells was lower. ** p




Anti-MTA1 antibody, PA1483, IHC(P)IHC(P): Human Rectal Cancer Tissue




IHC analysis of MTA1 using anti-MTA1 antibody (PA1483). MTA1 was detected in paraffin-embedded section of rat ovary tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-MTA1Antibody (PA1483) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.




Anti-MTA1 antibody, PA1483, IHC(F)IHC(F): Rat Ovary Tissue




IHC analysis of MTA1 using anti-MTA1 antibody (PA1483). MTA1 was detected in paraffin-embedded section of rat ovary tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-MTA1 Antibody (PA1483) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.




IHC analysis of MTA1 using anti-MTA1 antibody (PA1483). MTA1 was detected in frozen section of mouse brain tissues. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-MTA1 Antibody (PA1483) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.



IHC analysis of MTA1 using anti-MTA1 antibody (PA1483). MTA1 was detected in paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-MTA1 Antibody (PA1483) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.



IHC analysis of MTA1 using anti-MTA1 antibody (PA1483). MTA1 was detected in paraffin-embedded section of human ovarian cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-MTA1 Antibody (PA1483) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.



IHC analysis of MTA1 using anti-MTA1 antibody (PA1483). MTA1 was detected in paraffin-embedded section of human pancreas cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-MTA1 Antibody (PA1483) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-MTA1 Antibody

For Research Use Only. Not for use in diagnostic procedures.