

## Anti-Stra8 Antibody Picoband®

Catalog Number: PA1528

### About Stra8

Stra8, stimulated by retinoic acid 8, is expressed in embryonic ovaries just before meiotic initiation, whereas in testis it is expressed after birth. The STRA8 gene contains 9 exons. And this gene is mapped to chromosome 7q31. Retinoic acid (RA) signaling was required for Stra8 expression, and therefore initiation of meiosis, in embryonic mouse ovaries. RA was sufficient to induce Stra8 expression in embryonic testis. Cyp26-mediated RA metabolism was responsible for delaying Stra8 expression in embryonic testis. Since RA signaling regulated Stra8 expression in both embryonic ovaries and adult testis, it was said this portion of the meiotic initiation pathway may be identical in both sexes.

### Overview

Product Name	Anti-Stra8 Antibody Picoband®
Reactive Species	Rat
Description	Boster Bio Anti-Stra8 Antibody catalog # PA1528. Tested in WB applications. This antibody reacts with Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg Thimerosal, 0.05mg NaN <sub>3</sub> . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	D4ADQ4

### Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of rat Stra8, different from the related mouse sequence by one amino acid.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins

Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Rat

## Anti-Stra8 Antibody Picoband® (PA1528) Images



Anti-Stra8 antibody, PA1528, Western blotting  
All lanes: Anti Stra8(PA1528) at 0.5ug/ml  
WB : Rat Testis Tissue Lysate at 50ug  
Predicted bind size: 37KD  
Observed bind size: 37KD

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Stra8 Antibody

For Research Use Only. Not for use in diagnostic procedures.