

Anti-MCM2 Antibody Picoband® HRP Conjugated

Catalog Number: PA1711-HRP

About Mcm2

MCM2 (MINICHROMOSOME MAINTENANCE, S. CERVISIAE, HOMOLOG OF, 2), also known as MITOTIN, CDCL1 or BM28, is a human nuclear protein that plays an important role in 2 crucial steps of the cell cycle, namely, onset of DNA replication and cell division. And it is similar to members of the family of early S-phase proteins. The MCM2 gene is mapped to 3q21.3. The hexameric protein complex formed by MCM proteins is a key component of the pre-replication complex (pre-RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. In the G0 stage, the MCM2 and MCM5 proteins were much less abundant than the MCM7 and MCM3 proteins, which suggests that the MCM proteins are not present in stoichiometric amounts and that only a proportion of these molecules actively participate in cell cycle regulation as part of MCM complexes.

Overview

Product Name	Anti-MCM2 Antibody Picoband® HRP Conjugated
Reactive Species	Mouse, Rat
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	P97310

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of mouse MCM2, different from the related rat sequence by one amino acid.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	HRP
Suggested Dilutions	The intended application should be selected according to the customer's experimental requirements.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-MCM2 Antibody - HRP

For Research Use Only. Not for use in diagnostic procedures.