

## Anti-Replication factor C subunit 1 RFC1 Antibody Picoband® FITC Conjugated

Catalog Number: PA1716-FITC

### About RFC1

RFC (Replication Factor C), also called activator-1, is a multimeric primer-recognition protein consisting of 5 distinct subunits of 145, 40, 38, 37, and 36.5 kD. Human RFC was purified from extracts of HeLa cells as a host factor essential for the in vitro replication of simian virus 40 (SV40) DNA. RFC, in the presence of ATP, assembles proliferating-cell nuclear antigen and DNA polymerase-delta or polymerase-epsilon on primed DNA templates. Tight interfacial coordination of the ATP analog ATP-gamma-S by RFC resulted in a spiral arrangement of the ATPase domains of the clamp loader above the PCNA ring.

### Overview

Product Name	Anti-Replication factor C subunit 1 RFC1 Antibody Picoband® FITC Conjugated
Reactive Species	Human
Application	Flow Cytometry
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.02% NaN <sub>3</sub> .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P35251

### Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human RFC1.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	FITC Excitation Wavelength: 495 nm Emission Wavelength: 525 nm
Suggested Dilutions	Flow Cytometry, Optimal dilutions should be determined by end users.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Replication factor C subunit 1 RFC1 Antibody - FITC

For Research Use Only. Not for use in diagnostic procedures.