

Anti-CD2/CD2 Antibody Picoband® Fluoro594 Conjugated

Catalog Number: PA1743-Fluoro594

About Cd2

CD2 (cluster of differentiation 2) is a cell adhesion molecule found on the surface of T cells and natural killer (NK) cells. It has also been called T-cell surface antigen T11/Leu-5, LFA-2, LFA-3 receptor, erythrocyte receptor and rosette receptor. Monoclonal antibodies directed against CD2 inhibit the formation of rosettes with sheep erythrocytes, indicating that CD2 is the erythrocyte receptor or is closely associated with it. It is one of the earliest T-cell markers, being present on more than 95% of thymocytes; it is also found on some natural killer cells but not on B lymphocytes. Due to its structural characteristics, CD2 is a member of the immunoglobulin superfamily; it possesses two immunoglobulin-like domains in its extracellular portion. The localization of CD2 to 1p13 was established by in situ hybridization. By Southern blotting of DNA from a panel of somatic cell hybrids, Clayton et al. (1988) assigned the CD2 gene to human chromosome 1 and murine chromosome 3. CD2 interacts with other adhesion molecules, such as lymphocyte function-associated antigen-3 (LFA-3/CD58) in humans, or CD48 in rodents, which are expressed on the surfaces of other cells. With the use of transgenic mice, such an LCR was identified within the 3-prime flanking region of the human CD2 gene.

Overview

Product Name	Anti-CD2/CD2 Antibody Picoband® Fluoro594 Conjugated
Reactive Species	Mouse, Rat
Application	Flow Cytometry
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ N.
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P08920

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of mouse CD2, different from the related rat sequence by two amino acids.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.

Conjugate	Fluoro594 Excitation Wavelength: 593 nm Emission Wavelength: 618 nm
Suggested Dilutions	Flow Cytometry, Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CD2/CD2 Antibody - Fluoro594

For Research Use Only. Not for use in diagnostic procedures.