

Anti-CISH Antibody Picoband®

Catalog Number: PA1786

About CISH

CISH (cytokine inducible SH2-containing protein), also called CIS, CIS-1, G18, SOCS, is an important negative regulator for inflammatory signaling and belongs to the suppressors of cytokine signaling (SOCS) family. CIS family members are known to be cytokine-inducible negative regulators of cytokine signaling. CISH controls interleukin-2 signaling, and variations of CISH with certain SNPs are associated with susceptibility to bacteremia, tuberculosis and malaria. The human CISH gene is mapped to chromosome 3p21.3 by FISH. The mouse gene is tightly linked to the Gnai2 gene on chromosome 9, a region syntenic to human chromosome 3p21. CIS expression was upregulated by lipopolysaccharide (LPS) or *Cryptosporidium parvum* exposure, and this upregulation involved downregulation of MIR98 and LET7, which relieved MIR98- and LET7-mediated translational repression of CIS. Gain- and loss-of-function studies showed that CIS accelerated degradation of IKBA and enhanced NFκB activation in cholangiocytes in response to LPS stimulation or *C. parvum* exposure.

Overview


Product Name	Anti-CISH Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-CISH Antibody catalog # PA1786. Tested in IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9NSE2

Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human CISH, identical to the related rat and mouse sequences.
-----------	--

Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and ICC.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Immunocytochemistry , 0.5-1ug/ml, Human, Mouse, Rat Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, Rat, Mouse Western blot, 0.1-0.5ug/ml, Human, Rat, Mouse


Anti-CISH Antibody Picoband® (PA1786) Images



Western blot analysis of CISH using anti-CISH antibody (PA1786). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human MCF-7 whole cell lysates, Lane 2: human Hela whole cell lysates, Lane 3: human COLO320 whole cell lysates, Lane 4: human CACO-2 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CISH antigen affinity purified polyclonal antibody (Catalog # PA1786) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CISH at approximately 37 kDa. The expected band size for CISH is at 29 kDa.




IHC analysis of CISH using anti-CISH antibody (PA1786). CISH was detected in a paraffin-embedded section of Rat Kidney tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 ug/ml rabbit anti-CISH Antibody (PA1786) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IF analysis of CISH using anti-CISH antibody (PA1786). CISH was detected in an immunocytochemical section of U2OS cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The

IF analysis of CISH using anti-CISH antibody (PA1786). CISH was detected in an immunocytochemical section of U2OS cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The



cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-CISH Antibody (PA1786) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IF analysis of CISH using anti-CISH antibody (PA1786). CISH was detected in an immunocytochemical section of U2OS cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-CISH Antibody (PA1786) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CISH Antibody

For Research Use Only. Not for use in diagnostic procedures.