

Anti-TRIF/TICAM1 Antibody Picoband®

Catalog Number: PA1902

About TICAM1

TICAM (TIR DOMAIN-CONTAINING ADAPTOR MOLECULE 1), also known as TICAM1 or TRIF, is an adapter in responding to activation of toll-like receptors (TLRs). It mediates the rather delayed cascade of two TLR-associated signaling cascades, where the other one is dependent upon a MyD88 adapter. By genomic sequence analysis, Oshiumi et al. (2003) mapped the TICAM1 gene to chromosome 19p13.3. By coimmunoprecipitation analysis, Oshiumi et al. (2003) showed that TICAM1 interacts specifically with TLR3, but not with other TLRs. Functional analysis showed that the association of TLR3 and TICAM1 mediates dsRNA activation of IFN β , through either NF κ B, AP1, or IRF3. TICAM1 activation of NF κ B was found to occur predominantly through IRAK1 rather than IRAK2. Small interfering (si)RNA blockage of TICAM1, just upstream of the TIR domain, reduced IFN β production in response to dsRNA.

Overview


Product Name	Anti-TRIF/TICAM1 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-TRIF/TICAM1 Antibody catalog # PA1902. Tested in IHC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q8IUC6

Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human TRIF.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and IHC(F).

Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Immunohistochemistry (Frozen Section), 0.5-1ug/ml, Human, - Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human Western blot, 0.1-0.5ug/ml, Human


Anti-TRIF/TICAM1 Antibody Picoband® (PA1902) Images



Anti-TRIF antibody, PA1902, Western blotting
Recombinant Protein Detection Source: E.coli derived -recombinant Human TICAM1, 44.7KD(162aa tag+ Q468-E712)
Lane 1: Recombinant Human TICAM1 Protein 10ng
Lane 2: Recombinant Human TICAM1 Protein 5ng
Lane 3: Recombinant Human TICAM1 Protein 2.5ng
Lane 4: Recombinant Human TICAM1 Protein 1.25ng



Anti-TRIF antibody, PA1902, IHC(P)
IHC(P): Human Mammary Cancer Tissue



Western blot analysis of TRIF using anti-TRIF antibody (PA1902). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human Raji whole cell lysate. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TRIF antigen affinity purified polyclonal antibody (Catalog # PA1902) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TRIF at approximately 110KD. The expected band size for TRIF is at 76KD.

1 Publications Citing This Product

1. PubMed ID: 27164829, Curcumin exerts anti-inflammatory and antioxidative properties in 1-methyl-4-phenylpyridinium ion (MPP(+))-stimulated mesencephalic astrocytes by interference with TLR4 and downstream signaling pathway

Visit bosterbio.com/anti-trif-antibody-pa1902-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TRIF/TICAM1 Antibody

For Research Use Only. Not for use in diagnostic procedures.