

Anti-Macrosialin CD68 Antibody Picoband® FITC Conjugated

Catalog Number: PA2018-FITC

About CD68

CD68, cluster of differentiation, is a 110-kD transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. CD68 is a member of a family of hematopoietic mucin-like molecules that includes leukosialin/CD43 and stem cell antigen CD34. The CD68 gene is mapped to 17p13.1. Immunohistochemistry can be used to identify the presence of CD68, which is found in the cytoplasmic granules of a range of different blood cells. It is particularly useful as a marker for the various cells of the macrophage lineage, including monocytes, histiocytes, giant cells, Kupffer cells, and osteoclasts. This allows it to be used to distinguish diseases of otherwise similar appearance, such as the monocyte/macrophage and lymphoid forms of leukaemia (the latter being CD68 negative). Its presence in macrophages also makes it useful in diagnosing conditions related to proliferation or abnormality of these cells, such as malignant histiocytosis, histiocytic lymphoma, and Gaucher's disease.

Overview

Product Name	Anti-Macrosialin CD68 Antibody Picoband® FITC Conjugated
Reactive Species	Human
Application	Recommended applications are based on the parent unconjugated antibody (IHC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P34810

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human CD68.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	FITC Excitation Wavelength: 495 nm Emission Wavelength: 525 nm

Suggested Dilutions

Optimal dilutions should be determined by end users.

32 Publications Citing This Product

1. PubMed ID: -, Jian Pang, Li Zhou, Liqiu Liao et al. Correlation of sTILs Infiltration and UBR5 Expression on the Efficacy of Neoadjuvant Chemotherapy Efficacy and Prognosis in Early Triple Negative Breast Cancer, 13 January 2021, PREPRINT (Version 1) available at Research Square
2. PubMed ID: -, Fan Guo, Wei na Kong, Gang Zhao et al. The Correlation Between Tumor-associated Macrophage Infiltration and Progression in Cervical Carcinoma, 01 December 2020, PREPRINT (Version 1) available at Research Square[+https://doi.org/10.21203/rs.3.rs-115339/v1+]
3. PubMed ID: 32765086, Guo F, Feng YC, Zhao G, Zhang R, Cheng ZZ, Kong WN, Wu HL, Xu B, Lv X, Ma XM. Tumor-Associated CD163+ M2 Macrophage Infiltration is Highly Associated with PD-L1 Expression in Cervical Cancer. Cancer Manag Res. 2020 Jul 15;12:5831-5843. doi:10.2147/CMAR.S257692. PMID:32

Visit bosterbio.com/anti-cd68-antibody-pa2018-boster.html to see all 32 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Macrosialin CD68 Antibody - FITC

For Research Use Only. Not for use in diagnostic procedures.