

Anti-eIF3B Antibody Picoband®

Catalog Number: PA2030

About EIF3B

EIF3B (Eukaryotic Translation Initiation Factor 3, Subunit B), also called PRT1 or EIF3-p116, is a protein that in humans is encoded by the EIF3B gene. By searching an EST database, Methot et al. (1997) identified a cDNA encoding a human PRT1 homolog. Asano et al. (1997) demonstrated that the 115-kD component of HeLa cell eIF3 is actually composed of 2 proteins, p116 (PRT1) and an unrelated 110-kD protein. Chaudhuri et al. (1997) isolated cDNAs encoding PRT1, which they called p110 based on the size of the corresponding component of rabbit eIF3.

Overview


| | |
|----------------------|---|
| Product Name | Anti-eIF3B Antibody Picoband® |
| Reactive Species | Human |
| Description | Boster Bio Anti-eIF3B Antibody catalog # PA2030. Tested in WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance. |
| Application | WB |
| Clonality | Polyclonal |
| Formulation | Each vial contains antibody formulated with stabilizing components, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required. |
| Storage Instructions | Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles. |
| Host | Rabbit |
| Uniprot ID | P55884 |

Technical Details

| | |
|-------------------------------|---|
| Immunogen | A synthetic peptide corresponding to a sequence at the C-terminus of human eIF3B, different from the related rat and mouse sequences by one amino acid. |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot. |
| Cross Reactivity | No cross-reactivity with other proteins |
| Isotype | Rabbit IgG |

| | |
|---------------------|---|
| Form | Lyophilized |
| Concentration | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. |
| Purification | Immunogen affinity purified. |
| Suggested Dilutions | Western blot, 0.1-0.5ug/ml, Human |

Anti-eIF3B Antibody Picoband® (PA2030) Images



Western blot analysis of EIF3B using anti-EIF3B antibody (PA2030). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human A375 whole cell lysates, Lane 2: human K562 whole cell lysates, Lane 3: human MCF-7 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-EIF3B antigen affinity purified polyclonal antibody (Catalog # PA2030) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for EIF3B at approximately 115 kDa. The expected band size for EIF3B is at 92 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-eIF3B Antibody

For Research Use Only. Not for use in diagnostic procedures.