

888-466-3604 | support@bosterbio.com | www.bosterbio.com

# Anti-Bag5 Antibody

Catalog Number: PA2101

# About BAG5

BAG family molecular chaperone regulator 5 is a protein that in humans is encoded by the BAG5 gene. The protein encoded by this gene is a member of the BAG1-related protein family. Bag5 is a negative regulator of both Hsp70 and parkin function that sensitizes dopaminergic neurons to injury-induced death and thus promotes neurodegeneration.

### Overview

	1
Product Name	Anti-Bag5 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Bag5 Antibody catalog # PA2101. Tested in Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2mg Na2HPO4, 0.05 mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9UL15

## **Technical Details**

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human BAG5, different from the related mouse sequence by one amino acid, and from the related rat sequence by two amino acids.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.



888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat Flow Cytometry, 1-3 ug/1x10 <sup>6</sup> cells, Human



888-466-3604 | support@bosterbio.com | www.bosterbio.com

### Anti-Bag5 Antibody (PA2101) Images




Figure 1. Western blot analysis of BAG5 using anti-BAG5 antibody (PA2101).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Hela whole cell lysates, Lane 2: human HepG2 whole cell lysates,

Lane 3: rat testis tissue lysates,

Lane 4: mouse testis tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-BAG5 antigen affinity purified polyclonal antibody (Catalog # PA2101) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for BAG5 at approximately 51 kDa. The expected band size for BAG5 is at 51 kDa.




Figure 2. Flow Cytometry analysis of A431 cells using anti-BAG5 antibody (PA2101).

Overlay histogram showing A431 cells stained with PA2101 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-BAG5 Antibody (PA2101, 1 ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

#### Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Bag5 Antibody