

## Anti-Tyrosine-protein kinase Mer MERTK Antibody Picoband®

Catalog Number: PA2276

### About MERTK

Proto-oncogene tyrosine-protein kinase MER, also called MERTK is an enzyme that in humans is encoded by the MERTK gene. This gene is a member of the MER/AXL/TYRO3 receptor kinase family and encodes a transmembrane protein with two fibronectin type-III domains, two Ig-like C2-type (immunoglobulin-like) domains, and one tyrosine kinase domain. This gene is mapped to 2q13. MERTK signaling plays a role in various processes such as macrophage clearance of apoptotic cells, platelet aggregation, cytoskeleton reorganization and engulfment. This gene is also very important in inhibition of Toll-like receptors (TLRs)-mediated innate immune response by activating STAT1, which selectively induces production of suppressors of cytokine signaling SOCS1 and SOCS3.

### Overview


|                      |  |
|----------------------|--|
| Product Name         | Anti-Tyrosine-protein kinase Mer MERTK Antibody Picoband®  |
| Reactive Species     | Human, Monkey, Mouse, Rat  |
| Description          | Boster Bio Anti-Tyrosine-protein kinase Mer MERTK Antibody catalog # PA2276. Tested in Flow Cytometry, WB applications. This antibody reacts with Human, Monkey, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance. |
| Application          | Flow Cytometry, WB   |
| Clonality            | Polyclonal   |
| Formulation          | Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05 mg NaN <sub>3</sub> .  |
| Storage Instructions | Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.  |
| Host                 | Rabbit   |
| Uniprot ID           | Q12866   |

### Technical Details


|                               |   |
|-------------------------------|---|
| Immunogen                     | A synthetic peptide corresponding to a sequence at the N-terminus of human MERTK, different from the related mouse and rat sequences by one amino acid. |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.   |
| Cross Reactivity              | No cross-reactivity with other proteins   |
| Isotype                       | Rabbit IgG  |

|                     |  |
|---------------------|--|
| Form                | Lyophilized  |
| Concentration       | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.  |
| Purification        | Immunogen affinity purified.   |
| Suggested Dilutions | Western blot, 0.1-0.5ug/ml, Human, Monkey, Mouse, Rat<br>Flow Cytometry (Fixed), 1-3 ug/1x10 <sup>6</sup> cells, Human |

## Anti-Tyrosine-protein kinase Mer MERTK Antibody Picoband® (PA2276) Images



Western blot analysis of MERTK using anti-MERTK antibody (PA2276). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: human Jurkat whole cell lysates, Lane 3: human K562 whole cell lysates, Lane 4: monkey kidney tissue lysates, Lane 5: rat testis tissue lysates, Lane 6: mouse testis tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MERTK antigen affinity purified polyclonal antibody (Catalog # PA2276) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for MERTK at approximately 120 kDa. The expected band size for MERTK is at 110 kDa.



Flow Cytometry analysis of HepG2 cells using anti-MERTK antibody (PA2276). Overlay histogram showing HepG2 cells stained with PA2276 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-MERTK Antibody (PA2276, 1 ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

## 1 Publications Citing This Product

1. PubMed ID: 26621741, Functional screen identifies kinases driving prostate cancer visceral and bone metastasis

Visit [bosterbio.com/anti-mertk-antibody-pa2276-boster.html](http://bosterbio.com/anti-mertk-antibody-pa2276-boster.html) to see all 1 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Tyrosine-protein kinase Mer MERTK Antibody

For Research Use Only. Not for use in diagnostic procedures.