

Anti-ACTN3 Antibody Picoband®

Catalog Number: PB10026

About ACTN3

Alpha-actinin-3, also known as alpha-actinin skeletal muscle isoform 3 or F-actin cross-linking protein, is a protein that in humans is encoded by the ACTN3 gene. This gene encodes a member of the alpha-actin binding protein gene family. The encoded protein is primarily expressed in skeletal muscle and functions as a structural component of sarcomeric Z line. This protein is involved in crosslinking actin containing thin filaments. An allelic polymorphism in this gene results in both coding and non-coding variants; the reference genome represents the coding allele. The non-functional allele of this gene is associated with elite athlete status.

Overview


Product Name	Anti-ACTN3 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ACTN3 Antibody Picoband® catalog # PB10026. Tested in Flow Cytometry, IF, IHC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	Flow Cytometry, IF, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , and 0.05 mg NaN ₃ . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q08043

Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human ACTN3, different from the related mouse sequence by five amino acids.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).

Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, Mouse, Rat Immunofluorescence, 5 ug/ml, Mouse Flow Cytometry (Fixed), 1-3ug/1x10 ⁶ cells, Human


Anti-ACTN3 Antibody Picoband® (PB10026) Images



Western blot analysis of ACTN3/Alpha Actinin 3 using anti-ACTN3/Alpha Actinin 3 antibody (PB10026). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HT1080 whole cell lysates, Lane 2: rat skeletal muscle tissue lysates, Lane 3: rat skeletal muscle tissue lysates, Lane 4: mouse skeletal muscle tissue lysates, Lane 5: mouse skeletal muscle tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ACTN3/Alpha Actinin 3 antigen affinity purified polyclonal antibody (Catalog # PB10026) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for ACTN3/Alpha Actinin 3 at approximately 103 kDa. The expected band size for ACTN3/Alpha Actinin 3 is at 103 kDa.



IHC analysis of ACTN3/Alpha Actinin 3 using anti-ACTN3/Alpha Actinin 3 antibody (PB10026). ACTN3/Alpha Actinin 3 was detected in paraffin-embedded section of mouse skeletal muscle tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-ACTN3/Alpha Actinin 3 Antibody (PB10026) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.




Major patterns of protein distribution according to fiber type in human skeletal muscle. The fiber type distribution of representative proteins (indicated by gene name) is shown for each pattern, with values expressed as per cent of the maximal value. ACTN3 , alpha-actinin-3; CYCS , cytochrome c; MYL2 , MLC-2 slow; MYLK2 , myosin light chain kinase 2; RYR1 , ryanodine receptor 1; TNNC2 , fast troponin C Index in PubMed under a CC BY license. PMID: 34727990


Myofibrillar proteins: fiber type distribution of representative proteins. Proteins showing similar distribution in the different types are not shown in the figure. Values are expressed as per cent of the maximal value. ACTN2 , alpha-actinin-2; ACTN3 , alpha-actinin-3; CAPZA1 , CapZ alpha1; LMOD2 ,

Fiber type			1	2A	2X
Predominant in type 1 fibers					
100	0.3	0.3	TNNC1		
100	0.8	0.7	TNNI1		
100	42	39	TNNT1		
100	0.7	0.9	MYL2		
100	9	70	MYL3		
100	6	7	MYL6B		
100	25	9	MYOM3		
100	33	8	LRRC39		
100	28	26	LMO2		
100	78	66	ACTN2		
100	2	2	MYO22		
Predominant in type 2 fibers					
			TNNC2	1	91
			TNNI2	0.3	85
			TNNT3	0.6	100
			MYL1	24	100
			MYLFP	2	98
			MYBPC2	0.2	34
			MYO2	54	93
			ACTN3	0.9	28
			MYO23	45	52
			CAPZA1	17	16


leiomodlin-2; LRRC39, Myomasp; MYL1, MLC1/3-fast; MYL2, MLC-2slow; MYL3, MLC-1 slow; MYLPF, MLC2-fast; MYL6B, MLC-1sa; MYBPC1, myosin-binding protein C1; MYBPC2, myosin-binding protein C2; MYBPH, myosin-binding protein H; MYOM2, myomesin 2; MYOM3, myomesin 3; MYO22, myozenin 2; MYO23, myozenin 3; TNNC1, slow troponin C; TNNC2, fast troponin C; TNNI1, slow troponin I; TNNI2, fast troponin I; TNNT1, slow troponin T; TNNT2, fast troponin T; TPM1, alpha-tropomyosin; TPM3, gamma-tropomyosin
Index in PubMed under a CC BY license. PMID: 34727990



Fiber type-specific distribution of selected proteins revealed by immunofluorescence analysis. Fiber types are labeled as 1 (type 1), 2A (type 2A), or 2X (type 2X). Hybrid 2A-2X fibers are labeled by asterisks. A - D Type 1-specific proteins. Each panel shows on the left a section stained for PGM5/aciculin (A), PDLIM1 (B), MCU (C), or IDH2 (D) and on the right a serial section stained with MYH-specific antibodies to reveal the three fiber types. No pure type 2X fibers is present in A and B. E - F Type 2-specific proteins. Each panel shows on the left a section stained for ACTN3 (E) or XIRP2 (F), on the center a serial section stained for MYH-specific antibodies to reveal the three fiber types, and on the right a section stained with anti-MYH antibody BF-35, which reacts with MYH7 and MYH2 but not with MYH1, thus stains all fiber types, except pure type 2X fibers. Note that whereas ACTN3 is especially abundant in type 2X fibers, as well as in hybrid 2A-2X fibers, XIRP2 is present at higher levels in both 2A and 2X fibers. WGA counterstain was applied to all sections, except those processed with the three anti-MYHs antibodies
Index in PubMed under a CC BY license. PMID: 34727990




IHC analysis of ACTN3/Alpha Actinin 3 using anti-ACTN3/Alpha Actinin 3 antibody (PB10026). ACTN3/Alpha Actinin 3 was detected in paraffin-embedded section of rat skeletal muscle tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-ACTN3/Alpha Actinin 3 Antibody (PB10026) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.




IHC analysis of ACTN3/Alpha Actinin 3 using anti-ACTN3/Alpha Actinin 3 antibody (PB10026). ACTN3/Alpha Actinin 3 was detected in paraffin-embedded section of human lung cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-ACTN3/Alpha Actinin 3 Antibody (PB10026) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using

Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.



IF analysis of ACTN3/Alpha Actinin 3 using anti-ACTN3/Alpha Actinin 3 antibody (PB10026). ACTN3/Alpha Actinin 3 was detected in a paraffin-embedded section of mouse skeletal muscle tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/mL rabbit anti-ACTN3/Alpha Actinin 3 Antibody (PB10026) overnight at 4°C. FITC Conjugated Goat Anti-Rabbit IgG (BA1105) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



Flow Cytometry analysis of WISH cells using anti-ACTN3 antibody (PB10026). Overlay histogram showing WISH cells stained with PB10026 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ACTN3 Antibody (PB10026, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight® 488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

1 Publications Citing This Product

1. PubMed ID: 10.1186/s13395-021-00279-0, Protein profile of fiber types in human skeletal muscle: a single-fiber proteomics study

Visit bosterbio.com/anti-actn3-picoband-trade-antibody-pb10026-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ACTN3 Antibody

For Research Use Only. Not for use in diagnostic procedures.