

Anti-ADO Antibody Picoband® APC Conjugated

Catalog Number: PB10032-APC

About ADO

Human thiol dioxygenases include cysteine dioxygenase (CDO) and cysteamine (2-aminoethanethiol) dioxygenase (ADO). CDO adds 2 oxygen atoms to free cysteine, whereas ADO adds 2 oxygen atoms to free cysteamine to form hypotaurine. It is demonstrated that mouse Ado has strong and specific dioxygenase activity in vitro towards cysteamine but not cysteine. Recombinant Ado was shown to bind iron. Overexpression of Ado in HepG2/C3A cells increased the production of hypotaurine from cysteamine. Similar results were found with human ADO. When endogenous expression of ADO was reduced by RNA-mediated interference, hypotaurine production decreased. It is also noted that the demonstration of high levels of ADO in brain challenges the previous assumption that most of the taurine in the brain is a consequence of CDO activity.

Overview

Product Name	Anti-ADO Antibody Picoband® APC Conjugated
Reactive Species	Human, Mouse, Rat
Application	Flow Cytometry
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	Q96SZ5

Technical Details

Immunogen	E. coli-derived human ADO recombinant protein (Position: E49-E261). Human ADO shares 90.1% amino acid (aa) sequence identity with mouse ADO.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	APC Excitation Wavelength: 633-647 nm Emission Wavelength: 660 nm

Suggested Dilutions

Flow Cytometry, Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ADO Antibody - APC

For Research Use Only. Not for use in diagnostic procedures.