

Anti-APC2 Antibody Picoband®

Catalog Number: PB10041

About APC2

APC2, which is also called APCL, is a deduced 2,303-amino acid protein that contains an N-terminal coiled-coil domain, followed by an armadillo domain and five 20-amino acid repeats. The human APC2 gene is mapped to chromosome 19p13.3. It is found that the 20-amino acid repeat domain of APCL could bind beta-catenin (CTNNB1) and deplete the intracellular beta-catenin pool. A reporter gene assay revealed that APCL could regulate interaction of beta-catenin with T cell-specific transcription factors (TCF7), although less efficiently than APC.

Overview

Product Name	Anti-APC2 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-APC2 Antibody Picoband® catalog # PB10041. Tested in WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O95996

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human APC2, different from the related mouse sequence by two amino acids.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human

Anti-APC2 Antibody Picoband® (PB10041) Images

100KD — APC2
70KD —
55KD —
35KD —
25KD —
15KD —

Western blot analysis of APC2 using anti-APC2 antibody (PB10041). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: HELA whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-APC2 antigen affinity purified polyclonal antibody (Catalog # PB10041) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for APC2 at approximately 94 kDa. The expected band size for APC2 is at 94 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-APC2 Antibody

For Research Use Only. Not for use in diagnostic procedures.