

Anti-TNF alpha Antibody Picoband®

Catalog Number: PB9010

About Tnf

TNF alpha (Tumor Necrosis Factor alpha) gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFR2. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, and cancer. Knockout studies in mice also suggested the neuroprotective function of this cytokine.

Overview


Product Name	Anti-TNF alpha Antibody Picoband®
Reactive Species	Rat
Description	Boster Bio Anti-TNF alpha Antibody Picoband® catalog # PB9010. Tested in WB applications. This antibody reacts with Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P16599

Technical Details


Immunogen	E.coli-derived rat TNF alpha recombinant protein (Position: D89-L235). Rat TNF alpha shares 95% amino acid (aa) sequence identity with mouse TNF alpha.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Rat


Anti-TNF alpha Antibody Picoband® (PB9010) Images



Western blot analysis of TNF alpha using anti-TNF alpha antibody (PB9010). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: PC-12 Whole Cell Lysate, Lane 2: Rat Spleen Tissue Lysate, Lane 3: Rat Brain Tissue Lysate, Lane 4: Rat Kidney Tissue Lysate, Lane 5: Rat Liver Tissue Lysate. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TNF alpha antigen affinity purified polyclonal antibody (Catalog # PB9010) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TNF alpha at approximately 26KD. The expected band size for TNF alpha is at 26KD.




Red nucleus IL-33 facilitates the early development of mononeuropathic pain by inducing TNF-alpha through activating ERK, p38 MAPK and JAK2/STAT3 signaling pathways. A Western blotting showed that red nucleus TNF-alpha was increased at 1 week post-SNI, intrarubral injection of anti-IL-33 antibody at 1 week post-SNI suppressed the overexpression of TNF-alpha (n = 6 per group, F = 14.302, P < 0.001). B Western blotting displayed that intrarubral administration of PD98059, SB203580, or AG490 at 1 week post-SNI inhibited the production of TNF-alpha (n = 6 per group, F = 13.157, P < 0.001). C Immunohistochemistry showed that red nucleus TNF-alpha was upregulated at 1 week post-SNI, intrarubral injection of anti-IL-33 antibody, PD98059, SB203580, or AG490 at 1 week post-SNI suppressed the production of TNF-alpha (n = 4 per group, F = 33.029, P < 0.001). ** P < 0.01 and *** P < 0.001. Scale bars = 50 um Index in PubMed under a CC BY license. PMID: 34225736




Red nucleus IL-33 evokes mechanical hypersensitivity by inducing TNF-alpha through activating ERK, p38 MAPK, and JAK2/STAT3 signaling pathways. A Western blotting indicated that intrarubral administration of IL-33 stimulated the secretion of TNF-alpha in naive rats (n = 6 per group, F = 15.143, P < 0.001). B Western blotting showed that intrarubral pre-injection of PD98059, SB203580, or AG490, 30 min before IL-33 administration, restrained IL-33-induced overexpression of TNF-alpha in naive rats (n = 6 per group, F = 9.812, P < 0.001). C Immunohistochemistry showed that intrarubral injection of IL-33 potentiated the secretion of TNF-alpha in naive rats, intrarubral pre-injection of PD98059,


SB203580, or AG490, 30 min prior to IL-33 administration, inhibited IL-33-induced overexpression of TNF-alpha in naive rats (n = 4 per group, F = 44.310, P < 0.001). *** P < 0.001. Scale bars = 50 um Index in PubMed under a CC BY license. PMID: 34225736



DJ-1 interference increased the expression of TNF-alpha, IL-1beta, and IL-6 after cerebral I/R injury. a and c Western blot detecting DJ-1 and the cytokines TNF-alpha, IL-1beta, and IL-6 in rats. b, d Western blot detecting DJ-1 and the cytokines TNF-alpha, IL-1beta, and IL-6 in astrocytes. e-g Quantification of TNF-alpha, IL-1beta, and IL-6 in rats by ELISA. h-j Quantification of TNF-alpha, IL-1beta, and IL-6 in astrocytes by ELISA. The data are expressed as the mean ± SEM. * p < 0.05.




DJ-1 inhibited the expression of TNF-alpha, IL-1beta, and IL-6 after cerebral I/R injury via SHP-1. a, c After virus and TPI-1 were used to overexpress DJ-1 and inhibit SHP-1, respectively, Western blotting was used to detect the cytokines IL-1beta, IL-6, and TNF-alpha in rats. b, d After virus and TPI-1 were used to overexpress DJ-1 and inhibit SHP-1, respectively, Western blotting was used to detect the cytokines IL-1beta, IL-6, and TNF-alpha in astrocytes. e, g After treatment with an SHP-1 inhibitor, Western blotting was used to detect the cytokines IL-1beta, IL-6, and TNF-alpha in rats. f, h After treatment with an SHP-1 inhibitor, Western blotting was used to detect the cytokines IL-1beta, IL-6, and TNF-alpha in astrocytes. i, k, m Quantification of TNF-alpha, IL-1beta, and IL-6 in rats by ELISA. j, l, n Quantification of TNF-alpha, IL-1beta, and IL-6 in astrocytes by ELISA. The data are expressed as the mean ± SEM. * p < 0.05.




DJ-1 regulated the disassociation of NLRX1 from TRAF6 after cerebral I/R injury via SHP-1. a, g After virus and TPI-1 were used to overexpress DJ-1 and inhibit SHP-1, respectively, Western blotting was used to detect NLRX1, TRAF6, and SHP-1 in rats. b, h After virus and TPI-1 were used to overexpress DJ-1 and inhibit SHP-1, respectively, Western blotting was used to detect NLRX1, TRAF6, and SHP-1 in astrocytes. c, i After treatment with an SHP-1 inhibitor, Western blotting was used to detect the cytokines IL-1beta, IL-6, and TNF-alpha in rats. d, j After treatment with an SHP-1 inhibitor, Western blotting was used to detect the cytokines IL-1beta, IL-6, and TNF-alpha in astrocytes. e Immunoprecipitation and immunoblot analyses of NLRX1-TRAF6 in rats. f Immunoprecipitation and immunoblot analyses of SHP-1-TRAF6 in rats. The data are expressed as the mean ± SEM. * p < 0.05.

Fecal microbiota transplantation (FMT) reduced systemic or local inflammatory response. (A) Protein levels of the pro-inflammatory cytokines and chemokines in the blood. (B) mRNA levels of the pro-inflammatory cytokines and chemokines in kidney tissue using real-time PCR. (C)



Western blot analysis of the pro-inflammatory cytokines in kidney tissue. (D) Protein band intensities quantified using optical densitometry and normalized to beta-actin levels within the respective experimental groups. n = 6. Representative images from six independent experiments. The p -values above the data denote statistical comparisons on the groups—the control (Ctrl), ischemia-reperfusion injury plus normal saline (IRI+NS), and IRI+FMT groups—which were calculated using ANOVA with Tukey's post-hoc test. * p < 0.05; *** p < 0.001. Index in PubMed under a CC BY license. PMID: 41078364



Propionic acid reduced the systemic or local inflammatory response. (A) Protein levels of the pro-inflammatory cytokines and chemokines in the blood. (B) mRNA levels of the pro-inflammatory cytokines and chemokines in kidney tissue using real-time PCR. (C) Western blot analysis of the pro-inflammatory cytokines in kidney tissue. (D) Protein band intensities quantified using optical densitometry and normalized to beta-actin levels within the respective experimental groups. n = 6. Representative images from six independent experiments. The p -values above the data denote statistical comparisons of the groups—the control (Ctrl), ischemia-reperfusion injury plus normal saline (IRI+NS), and IRI plus propionic acid (IRI+Prop) groups—which were calculated using ANOVA with Tukey's post-hoc test. ** p < 0.01; *** p < 0.001. Index in PubMed under a CC BY license. PMID: 41078364

78 Publications Citing This Product

1. PubMed ID: 34225736, Li HN, Yang QQ, Wang WT, Tian X, Feng F, Zhang ST, Xia YT, Wang JX, Zou YW, Wang JY, Zeng XY. Red nucleus IL-33 facilitates the early development of mononeuropathic pain in male rats by inducing TNF-alpha through activating ERK, p38 MAPK, and JAK2/STAT3. *J Neuroinflammation*. 2021 Jul 5;18(1):150. doi:10.1186/s12974-021-02198-9. PMID:34225736; PMCID:PMC8258957.
2. PubMed ID: 33823825, Wang B, Gao C, Zhang P, Sun W, Zhang J, Gao J. The increased motion of lumbar induces ligamentum flavum hypertrophy in a rat model. *BMC Musculoskelet Disord*. 2021 Apr 6;22(1):334. doi:10.1186/s12891-021-04203-x. PMID:33823825.
3. PubMed ID: -, Yang Ping, Yingpeng Li, Shaowa Lü, Yali Sun, Wanmeng Zhang, Jialin Wu, Ting Liu, Yongji Li. A study of nanometre aggregates formation mechanism and antipyretic effect in Bai-Hu-Tang, an ancient Chinese herbal decoction. *Biomedicine & Pharmacotherapy*, Volume 124, 202

Visit bosterbio.com/anti-tnf-alpha-picoband-trade-antibody-pb9010-boster.html to see all 78 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TNF alpha Antibody

For Research Use Only. Not for use in diagnostic procedures.