

Anti-IL2 Antibody Picoband®

Catalog Number: PB9013

About IL2

Interleukin-2 (IL2), formerly referred to as T-cell growth factor, is a powerfully immunoregulatory lymphokine that is produced by lectin- or antigen-activated T cells. It is produced not only by mature T lymphocytes on stimulation but also constitutively by certain T-cell lymphoma cell lines. The lymphokine interleukin-2 (IL-2) is responsible for autocrine cell cycle progression and regulation of immune responses. IL-2 expression in mature thymocytes and T cells has been found to be tightly controlled by monoallelic expression. IL-2 can act as a growth hormone for both B and T lymphocytes. This gene for interleukin 2 (IL2) is assigned to chromosome 4. IL-2 is a 15.5KDa glycoprotein, consisting of 153 amino acids in precursor form and 133 amino acids in mature form.

Overview


Product Name	Anti-IL2 Antibody Picoband®
Reactive Species	Rat
Description	Boster Bio Anti-IL2 Antibody Picoband® catalog # PB9013. Tested in WB applications. This antibody reacts with Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , and 0.05 mg NaN ₃ . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P17108

Technical Details

Immunogen	E.coli-derived rat IL-2 recombinant protein (Position: A21-Q155). Rat IL-2 shares 66% and 80% amino acid (aa) sequences identity with human and mouse IL-2, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western

	blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Rat

Anti-IL2 Antibody Picoband® (PB9013) Images



Western blot analysis of IL2 using anti-IL2 antibody (PB9013). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: Rat Thymus Tissue Lysate, Lane 2: Rat Liver Tissue Lysate, Lane 3: NRK Whole Cell Lysate, Lane 4: PC12 Whole Cell Lysate, Lane 5: RH35 Whole Cell Lysate. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IL2 antigen affinity purified polyclonal antibody (Catalog # PB9013) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for IL2 at approximately 18KD. The expected band size for IL2 is at 18KD.

7 Publications Citing This Product

1. PubMed ID: 25667679, Chen B, Liu L, Xu H, Yang Y, Zhang L, Zhang F. Exp Ther Med. 2015 Mar;9(3):1063-1067. Epub 2015 Jan 26. Effectiveness Of Immune Therapy Combined With Chemotherapy On The Immune Function And Recurrence Rate Of Cervical Cancer.
2. PubMed ID: 22594905, Zhong B, Shen H, Sun X, Wang H, Zhang Y, Sun Z. Cancer Biother Radiopharm. 2012 May;27(4):252-8. Doi: 10.1089/Cbr.2011.1105. Additive Effects Of Ulinastatin And Docetaxel On Growth Of Breast Cancer Xenograft In Nude Mice And Expression Of Pge2, Il...
3. PubMed ID: 28902929, Women with Recurrent Miscarriage Have Decreased Expression of 25-Hydroxyvitamin D3-1?-Hydroxylase by the Fetal-Maternal Interface

Visit bosterbio.com/anti-il-2-picoband-trade-antibody-pb9013-boster.html to see all 7 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-IL2 Antibody

For Research Use Only. Not for use in diagnostic procedures.