

## Anti-Interferon gamma/IFNG Antibody Picoband™

Catalog Number: PB9024

### About Ifng

Interferon-gamma (IFN-gamma) is an inflammatory cytokine that has been implicated in the development of fibrosis in inflamed tissues. The production of IFN-gamma, which is under genetic control, can influence the development of fibrosis in lung allografts. IFN-gamma is also produced by natural killer (NK) cells and most prominently by CD8 cytotoxic T cells, and is vital for the control of microbial pathogens. Interferon gamma is believed to be crucial for host defence against many infections. Genetically determined variability in IFN-gamma and expression might be important for the development of tuberculosis. IFN-gamma activates human macrophage oxidative metabolism and antimicrobial activity. In addition to having antiviral activity, IFN-gamma has important immunoregulatory functions. IFN-gamma plays an important role in the control of neointima proliferation.

### Overview

Product Name	Anti-Interferon gamma/IFNG Antibody Picoband™
Reactive Species	Rat
Description	Boster Bio Anti-Interferon gamma/IFNG Antibody Picoband™ catalog # PB9024. Tested in ELISA, WB applications. This antibody reacts with Rat.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P01581

### Technical Details

Immunogen	E.coli-derived rat IFN gamma recombinant protein (Position: Q23-C156). Rat IFN gamma shares 38% and 86% amino acid (aa) sequences identity with human and mouse IFN gamma, respectively.
Predicted Reactive Species	Bovine, Chicken, Horse, Monkey, Rabbit
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>ELISA , 0.1-0.5ug/ml, Rat, -</p> <p>Western blot, 0.1-0.5ug/ml, Rat</p>

## Anti-Interferon gamma/IFNG Antibody Picoband™ (PB9024) Images




Figure 1. Western blot analysis of Interferon gamma using anti-Interferon gamma antibody (PB9024).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours.

Lane 1: Recombinant Rat Interferon gamma Protein 0.5ng.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Interferon gamma antigen affinity purified polyclonal antibody (Catalog # PB9024) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Interferon gamma at approximately 19KD. The expected band size for Interferon gamma is at 19KD.

## 5 Publications Citing This Product

1. PubMed ID: 26222793, Somatostatin Improved B Cells Mature in Macaques during Intestinal Ischemia-Reperfusion
2. PubMed ID: 28902929, Women with Recurrent Miscarriage Have Decreased Expression of 25-Hydroxyvitamin D3-1?-Hydroxylase by the Fetal-Maternal Interface
3. PubMed ID: 25483698, Yan X, Wang D, Liang F, Fu L, Guo C. Hum Vaccin Immunother. 2014;10(12):3491-8. Doi: 10.4161/Hv.36084. Hpv16L1-Attenuated Shigella Recombinant Vaccine Induced Strong Vaginal And Systemic Immune Responses In Guinea Pig Model.

Visit [bosterbio.com/anti-ifn-gamma-picoband-trade-antibody-pb9024-boster.html](http://bosterbio.com/anti-ifn-gamma-picoband-trade-antibody-pb9024-boster.html) to see all 5 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Interferon gamma/IFNG Antibody™