

Anti-TARC/CCL17 Antibody Picoband® Fluoro488 Conjugated

Catalog Number: PB9031-Fluoro488

About Ccl17

Chemokine (C-C motif) ligand 17 (CCL17) is a small cytokine belonging to the CC chemokine family that is also known as thymus and activation regulated chemokine (TARC). CCL17 is expressed constitutively in thymus, but only transiently in phytohemagglutinin-stimulated peripheral blood mononuclear cells. This chemokine specifically binds and induces chemotaxis in T cells and elicits its effects by interacting with the chemokine receptor CCR4. The gene for CCL17 is located on chromosome 16, in humans, along with other chemokines called CCL22 and CX3CL1. The standard used in this kit is recombinant human CCL17, consisting of 71 amino acids with the molecular weight of 8Kda.

Overview

Product Name	Anti-TARC/CCL17 Antibody Picoband® Fluoro488 Conjugated
Reactive Species	Mouse
Application	Flow Cytometry
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	Q9WUZ6

Technical Details

Immunogen	E.coli-derived mouse TARC recombinant protein (Position: A34-P103). Mouse TARC shares 71% amino acid (aa) sequence identity with human TARC.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	Fluoro488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Suggested Dilutions	Flow Cytometry, Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TARC/CCL17 Antibody - Fluoro488

For Research Use Only. Not for use in diagnostic procedures.