

Anti-ARA9/AIP Antibody Picoband®

Catalog Number: PB9042

About AIP

AIP, also known as, ARA9 or XAP-2, is a protein that in humans is encoded by the AIP gene. This gene is mapped to 11q13.2. The encoded protein is found in the cytoplasm as part of a multiprotein complex, but upon binding of ligand is transported to the nucleus. AIP may play a positive role in aryl hydrocarbon receptor-mediated signalling possibly by influencing its receptivity for ligand and/or its nuclear targeting. It has been shown that AIP is the cellular negative regulator of the hepatitis B virus (HBV) X protein. AIP mutations may be the cause of a familial form of acromegaly, familial isolated pituitary adenoma (FIPA).

Overview


Product Name	Anti-ARA9/AIP Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-ARA9/AIP Antibody Picoband® catalog # PB9042. Tested in Flow Cytometry, IF, ICC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	Flow Cytometry, IF, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , and 0.05 mg NaN ₃ . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O00170

Technical Details


Immunogen	E.coli-derived human ARA9 recombinant protein (Position: D91-H330). Human ARA9 shares 95% amino acid (aa) sequence identity with both mouse and rat ARA9.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for ICC.

Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human Immunocytochemistry/Immunofluorescence, 2ug/ml, Human Flow Cytometry (Fixed), 1-3ug/1x10 ⁶ cells, Human


Anti-ARA9/AIP Antibody Picoband® (PB9042) Images



Western blot analysis of ARA9 using anti-ARA9 antibody (PB9042). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Hela whole cell lysates, Lane 2: human COLO320 whole cell lysates, Lane 3: human MCF-7 whole cell lysates, Lane 4: human SW620 whole cell lysates, Lane 5: human U87 whole cell lysates, Lane 6: human MDA-MB-453 whole cell lysates, Lane 7: human Jurkat whole cell lysates, Lane 8: human CACO-2 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ARA9 antigen affinity purified polyclonal antibody (Catalog # PB9042) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for ARA9 at approximately 38 kDa. The expected band size for ARA9 is at 38 kDa.



IF analysis of ARA9 using anti-ARA9 antibody (PB9042). ARA9 was detected in immunocytochemical section of U2OS cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 2ug/mL rabbit anti-ARA9 Antibody (PB9042) overnight at 4°C. DyLight@488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



Flow Cytometry analysis of A431 cells using anti-ARA9 antibody (PB9042). Overlay histogram showing A431 cells stained with PB9042 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ARA9 Antibody (PB9042, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight@488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ARA9/AIP Antibody

For Research Use Only. Not for use in diagnostic procedures.