BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-GDNF Antibody Picoband®

Catalog Number: PB9069

About GDNF

Glial cell line-derived neurotrophic factor (GDNF) is a glycosylated, disulfide-bonded homodimer that is a distantly related member of the transforming growth factor-beta superfamily. GDNF is also a potent neurotrophic factor that promotes the survival of dopaminergic neurones in cultures including embryonic neuronal cultures. In addition to its potential role in the differentiation and survival of central nervous system neurons, it has profound effects on kidney organogenesis and the development of the peripheral nervous system. GDNF may have utility in the treatment of Parkinson's disease, which is marked by progressive degeneration of midbrain dopaminergic neurons.

Submit a product review to Biocompare.com



Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-GDNF Antibody Picoband®
Reactive Species	Human, Rat
Description	Boster Bio Anti-GDNF Antibody Picoband® catalog # PB9069. Tested in WB applications. This antibody reacts with Human, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na2HPO4, and 0.05 mg NaN3. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20˚C for one year from date of receipt. After reconstitution, at 4˚C for one month. It can also be aliquotted and stored frozen at -20˚C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P39905

Technical Details

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Immunogen	E.coli-derived human GDNF recombinant protein (Position: S78-I211). Human GDNF shares 93% amino acid (aa) sequence identity with both mouse and rat GDNF.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Rat, Human

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-GDNF Antibody Picoband® (PB9069) Images



Western blot analysis of GDNF using anti-GDNF antibody (PB9069). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat brain tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-GDNF antigen affinity purified polyclonal antibody (Catalog # PB9069) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for GDNF at approximately 24 kDa. The expected band size for GDNF is at 24 kDa.

5 Publications Citing This Product

1. PubMed ID: 24503145, Zhang X, Shi K, Li Y, Zhang H, Hao J. Reprod Toxicol. 2014 Jun;45:87-93. Doi: 10.1016/J.Reprotox.2014.01.009. Epub 2014 Feb 4. Lipopolysaccharide Inhibits The Self-Renewal Of Spermatogonial Stem Cells In Vitro Via Downregulation Of Gdnf Expression...

2. PubMed ID: 21738758, Li R, Xia W, Zhang Z, Wu K. Plos One. 2011;6(6):E21663. Doi: 10.1371/Journal.Pone.0021663. Epub 2011 Jun 27. S100B Protein, Brain-Derived Neurotrophic Factor, And Glial Cell Line-Derived Neurotrophic Factor In Human Milk.

3. PubMed ID: 21738758, Li R, Xia W, Zhang Z, Wu K. Plos One. 2011;6(6):E21663. Doi: 10.1371/Journal.Pone.0021663. Epub 2011 Jun 27. S100B Protein, Brain-Derived Neurotrophic Factor, And Glial Cell Line-Derived Neurotrophic Factor In Human Milk.

Visit bosterbio.com/anti-gdnf-picoband-trade-antibody-pb9069-boster.html to see all 5 publications.

For Research Use Only. Not for use in diagnostic procedures.