

## Anti-Heme Oxygenase 1/HMOX1 Antibody Picoband®

Catalog Number: PB9085

### About Hmox1

HMOX1 (heme oxygenase (decycling) 1) is a human gene that encodes for the enzyme heme oxygenase 1. It is an essential enzyme in heme catabolism, it cleaves heme to form biliverdin. HMOX1 belongs to the heme oxygenase family. The HMOX1 gene is located on the long (q) arm of chromosome 22 at position 12.3, from base pair 34,101,636 to base pair 34,114,748. HMOX1, an essential enzyme in heme catabolism, cleaves heme to form biliverdin, which is subsequently converted to bilirubin by biliverdin reductase, and carbon monoxide, a putative neurotransmitter. HMOX1 activity is induced by its substrate heme and by various nonheme substances.

### Overview


|                      |                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|----------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Name         | Anti-Heme Oxygenase 1/HMOX1 Antibody Picoband®                                                                                                                                                                                                                                                                                                                                                                                      |
| Reactive Species     | Mouse, Rat                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Description          | Boster Bio Anti-Heme Oxygenase 1/HMOX1 Antibody Picoband® catalog # PB9085. Tested in IHC, WB applications. This antibody reacts with Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance. |
| Application          | IHC, WB                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Clonality            | Polyclonal                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Formulation          | Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .                                                                                                                                                                                                                                                                                                                                        |
| Storage Instructions | At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.                                                                                                                                                                                                                                   |
| Host                 | Rabbit                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Uniprot ID           | P14901                                                                                                                                                                                                                                                                                                                                                                                                                              |

### Technical Details


|                               |                                                                                                                                                                                  |
|-------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Immunogen                     | E.coli-derived mouse HMOX1 recombinant protein (Position: E2-T261). Mouse HMOX1 shares 82% and 93% amino acid (aa) sequences identity with human and rat HMOX1, respectively.    |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P). |
| Cross Reactivity              | No cross-reactivity with other proteins                                                                                                                                          |
| Isotype                       | Rabbit IgG                                                                                                                                                                       |
| Form                          | Lyophilized                                                                                                                                                                      |

|                     |                                                                                                                   |
|---------------------|-------------------------------------------------------------------------------------------------------------------|
| Concentration       | Add 0.2ml of distilled water will yield a concentration of 500ug/ml.                                              |
| Purification        | Immunogen affinity purified.                                                                                      |
| Suggested Dilutions | Western blot, 0.25-0.5ug/ml, Mouse, Rat<br>Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Mouse, Rat |


## Anti-Heme Oxygenase 1/HMOX1 Antibody Picoband® (PB9085) Images




Effects of magnolol on mice alcohol-induced liver damage in the Nrf2/HO-1 signaling pathway. After the completion of modeling and samples were collected, the liver of mice was lysed to detect the proteins by western blotting analysis. The levels of Nrf2 and HO-1 were compared with GAPDH. The data were demonstrated as means  $\pm$  SD. (\* $P < 0.05$ , \*\*\* $P < 0.001$  and "ns" means not significant). Index in PubMed under a CC BY license. PMID: 31920652



Trimetazidine restrained EE-induced activation of NF-kappaB and inactivation of HO-1/Nrf2 signaling pathways. (A-C) Western blot analysis of the protein levels of nuclear factor kappa light polypeptide gene enhancer in B-cells inhibitor alpha (IkappaBalpha), P-IkappaBalpha, and nuclear factor kappa-light-chain-enhancer of activated B-cells (NF-kappaB) in myocardial tissues. GAPDH and Histone H3 were used as loading controls. (D,E) The DNA binding activity of NF-kappaB was evaluated by EMSA assay. (F-G) Western blot analysis of the protein levels of heme oxygenase 1 (HO-1) and nuclear factor (erythroid-derived 2)-like 2 (Nrf2) in myocardial tissues. Each value is shown as mean  $\pm$  SD (n = 6). \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ , versus the indicated group. Index in PubMed under a CC BY license. PMID: 30890937




Involvement of Nrf2/HO-1 signaling pathway in therapeutic effects of NMN. (A) Representative images and quantitative analysis of HO-1 protein in peri-hemorrhagic brain tissues at 24 hours after cICH. (B) Representative images and quantitative analysis of Nrf2 protein in peri-hemorrhagic brain tissues at 24 hours after cICH. (C) Representative images and quantitative analysis of Nrf2 protein distribution in cytosolic and nuclear extracts of brain peri-hemorrhagic tissues at 24 hours after cICH. \*  $P < 0.05$




Western blot analysis of HMOX1 using anti-HMOX1 antibody (PB9085). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30  $\mu$ g of sample under reducing conditions. Lane 1: rat spleen tissue lysates, Lane 2: rat liver tissue lysates, Lane 3: rat PC-12 whole cell lysates, Lane 4: mouse spleen tissue lysates, Lane 5: mouse NIH/3T3 whole cell lysates, Lane 6: mouse RAW264.7 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-HMOX1 antigen affinity purified polyclonal antibody (PB9085) at 0.5  $\mu$ g/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary

antibody (Catalog # BA1054) at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for HMOX1 at approximately 33 kDa. The expected band size for HMOX1 is at 33 kDa.



IHC analysis of HMOX1 using anti-HMOX1 antibody (PB9085). HMOX1 was detected in a paraffin-embedded section of mouse liver tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-HMOX1 Antibody (PB9085) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of HMOX1 using anti-HMOX1 antibody (PB9085). HMOX1 was detected in a paraffin-embedded section of rat alcoholic hepatitis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-HMOX1 Antibody (PB9085) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-Heme Oxygenase 1/HMOX1 Antibody

For Research Use Only. Not for use in diagnostic procedures.