

Anti-Cyclin D3/CCND3 Antibody Picoband®

Catalog Number: PB9103

About CCND3

CCND3, also called Cyclin D3, is a protein that in humans is encoded by the CCND3 gene. It is mapped to 6p21.1. The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. In addition, overexpression of CCND3 upregulated the translational activity in HeLa cells in a dose-dependent manner.

Overview


Product Name	Anti-Cyclin D3/CCND3 Antibody Picoband®
Reactive Species	Human, Rat
Description	Boster Bio Anti-Cyclin D3/CCND3 Antibody Picoband® catalog # PB9103. Tested in WB applications. This antibody reacts with Human, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9mg NaCl, 0.2mg Na2HPO4, 0.01mg NaN3. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P30281

Technical Details


Immunogen	E.coli-derived human Cyclin D3 recombinant protein (Position: Q136-L292). Human Cyclin D3 shares 94% and 93% amino acid (aa) sequences identity with mouse and rat Cyclin D3, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western

	blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human, Rat

Anti-Cyclin D3/CCND3 Antibody Picoband® (PB9103) Images



Western blot analysis of Cyclin D3 using anti-Cyclin D3 antibody (PB9103). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. lane 1: recombinant human Cyclin D3 protein 0.5ng. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Cyclin D3 antigen affinity purified polyclonal antibody (Catalog # PB9103) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Cyclin D3 at approximately 25KD. The expected band size for Cyclin D3 is at 25KD.



Western blot analysis of Cyclin D4 using anti-Cyclin D4 antibody (PB9103). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. lane 1: rat testis tissue lysate, lane 2: rat thymus tissue lysate, lane 3: rat lung tissue lysate, lane 4: rat ovary tissue lysate, lane 5: JURKAT whole cell lysate, lane 6: A549 whole cell lysate, lane 7: MCF-7 whole cell lysate, lane 8: HELA whole cell lysate. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Cyclin D4 antigen affinity purified polyclonal antibody (Catalog # PB9103) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Cyclin D4 at approximately 33KD. The expected band size for Cyclin D4 is at 33KD.

1 Publications Citing This Product

1. PubMed ID: 10.1007/s10616-018-0235-3, Co-culture with TM4 cells enhances the proliferation and migration of rat adipose-derived mesenchymal stem cells with high stemness

Visit bosterbio.com/anti-cyclin-d3-picoband-trade-antibody-pb9103-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Cyclin D3/CCND3 Antibody

For Research Use Only. Not for use in diagnostic procedures.