

## Anti-EPB41L1 Antibody Picoband™

Catalog Number: PB9179

### About EPB41L1

Band 4.1-like protein 1 is a protein that in humans is encoded by the EPB41L1 gene. This gene is mapped to 20q11.23. It is found that a heterozygous missense mutation in the EPB41L1 gene causing nonsyndromic intellectual disability. This gene may function to confer stability and plasticity to neuronal membrane via multiple interactions, including the spectrin-actin-based cytoskeleton, integral membrane channels and membrane-associated guanylate kinases. What's more, EPB41L1 is a multifunctional protein that mediates interactions between the erythrocyte cytoskeleton and the overlying plasma membrane.

### Overview

|                      |   |
|----------------------|---|
| Product Name         | Anti-EPB41L1 Antibody Picoband™   |
| Reactive Species     | Human, Mouse, Rat   |
| Description          | Boster Bio Anti-EPB41L1 Antibody Picoband™ catalog # PB9179. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.  |
| Application          | WB  |
| Clonality            | Polyclonal  |
| Formulation          | Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.   |
| Storage Instructions | Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles. |
| Host                 | Rabbit  |
| Uniprot ID           | Q9H4G0  |

### Technical Details

|                               |   |
|-------------------------------|---|
| Immunogen                     | E.coli-derived human EPB41L1 recombinant protein (Position: Q691-S881). Human EPB41L1 shares 91% amino acid (aa) sequence identity with both mouse and rat EPB41L1. |
| Predicted Reactive Species    | Hamster   |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.   |
| Cross Reactivity              | No cross-reactivity with other proteins   |
| Isotype                       | Rabbit IgG  |
| Form                          | Lyophilized   |
| Concentration                 | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.   |

|                     |  |
|---------------------|--|
| Purification        | Immunogen affinity purified.   |
| Suggested Dilutions | <p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat</p> |

## Anti-EPB41L1 Antibody Picoband™ (PB9179) Images



**Figure 1. Western blot analysis of EPB41L1 using anti-EPB41L1 antibody (PB9179).**  
Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. Lane 1: recombinant human EPB41L1 protein 0.5 ng. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-EPB41L1 antigen affinity purified polyclonal antibody (Catalog # PB9179) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for EPB41L1 at approximately 50 kDa. The expected band size for EPB41L1 is at 50 kDa.



**Figure 2. Western blot analysis of EPB41L1 using anti-EPB41L1 antibody (PB9179).**  
Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human rat brain tissue lysates, Lane 2: mouse brain tissue lysates, Lane 3: human A549 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-EPB41L1 antigen affinity purified polyclonal antibody (Catalog # PB9179) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for EPB41L1 at approximately 98 kDa. The expected band size for EPB41L1 is at 99 kDa.

**Submit a product review to Biocompare.com**

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

