

Anti-Cytochrome P450 2E1/CYP2E1 Antibody Picoband® PE Conjugated

Catalog Number: PB9190-PE

About CYP2E1

Cytochrome P450 2E1 (abbreviated CYP2E1), a member of the cytochrome P450 mixed-function oxidase system, is involved in the metabolism of xenobiotics in the body. In humans, the CYP2E1 enzyme is encoded by the CYP2E1 gene. It is mapped to 10q26.3. While it is involved in the oxidative metabolism of a small range of substrates (mostly small polar molecules), there are many important drug interactions mediated by CYP2E1. Most drugs undergo deactivation by CYP2E1, either directly or by facilitated excretion from the body. Also, many substances are bioactivated by CYP2E1 to form their active compounds. In addition, CYP2E1 is an important enzyme for the conversion of ethanol to acetaldehyde and to acetate in humans. In the conversion sequence of acetyl-CoA to glucose, CYP2E1 transforms acetone via acetol into propylene glycol and methylglyoxal, the precursors of pyruvate, acetate and lactate.

Overview

| | |
|----------------------|---|
| Product Name | Anti-Cytochrome P450 2E1/CYP2E1 Antibody Picoband® PE Conjugated |
| Reactive Species | Mouse, Rat |
| Application | Recommended applications are based on the parent unconjugated antibody (IHC, WB). Customers may select suitable applications according to their experimental needs. |
| Clonality | Polyclonal |
| Formulation | Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% Na ₃ N. |
| Storage Instructions | At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light. |
| Host | Rabbit |
| Uniprot ID | P05181 |

Technical Details

| | |
|------------------|--|
| Immunogen | E.coli-derived human CYP2E1 recombinant protein (Position: M1-Y310). Human CYP2E1 shares 73% and 74% amino acid (aa) sequences identity with mouse and rat CYP2E1, respectively. |
| Cross Reactivity | No cross-reactivity with other proteins |
| Isotype | Rabbit IgG |
| Form | Liquid |
| Concentration | 0.5 mg/mL |
| Purification | Immunogen affinity purified. |
| Conjugate | PE Excitation Wavelength: 566 nm |

| | |
|---------------------|--|
| | Emission Wavelength: 574 nm |
| Suggested Dilutions | Optimal dilutions should be determined by end users. |

7 Publications Citing This Product

1. PubMed ID: 10.1002/ptr.5867, Nephroprotective Effects of Anthocyanin from the Fruits of Panax ginseng (GFA) on Cisplatin-Induced Acute Kidney Injury in Mice
2. PubMed ID: -, Thida,M.,Li,B.,Zhang,X.,Chen,C.,& Zhang,X.(2021).Echinacoside alleviates acetaminophen-induced liver injury by attenuating oxidative stress and inflammatory cytokines in mice.J Appl Biomed,19(2),112.doi:10.32725/jab.2021.011
3. PubMed ID: 33354276, Zhai T,Xu W,Liu Y,Qian K,Xiong Y,Chen Y.Honokiol Alleviates Methionine-Choline Deficient Diet-Induced Hepatic Steatosis and Oxidative Stress in C57BL/6 Mice by Regulating CFLAR-JNK Pathway.Oxid Med Cell Longev.2020 Nov 27;2020:2313641.doi:10.1155/2020/231

Visit bosterbio.com/anti-cyp2e1-picoband-trade-antibody-pb9190-boster.html to see all 7 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Cytochrome P450 2E1/CYP2E1 Antibody - PE

For Research Use Only. Not for use in diagnostic procedures.