

## Anti-Estrogen Receptor/ESR1 Antibody Picoband™

Catalog Number: PB9192

### About ESR1

Estrogen receptor alpha (ER-alpha), also known as NR3A1, is one of two main types of estrogen receptor, a nuclear receptor that is activated by the sex hormone estrogen. Estrogen receptors are involved in pathological processes including breast cancer, endometrial cancer, and osteoporosis. In humans, ER-alpha is encoded by the gene ESR1 (Estrogen Receptor 1). It is mapped to 6q25.1. This gene is a ligand-activated transcription factor composed of several domains important for hormone binding, DNA binding, and activation of transcription. The protein localizes to the nucleus where it may form a homodimer or a heterodimer with estrogen receptor 2. Estrogen and its receptors are essential for sexual development and reproductive function, it also play a role in other tissues such as bone.

### Overview

Product Name	Anti-Estrogen Receptor/ESR1 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-Estrogen Receptor/ESR1 Antibody Picoband™ catalog # PB9192. Tested in IHC, WB applications. This antibody reacts with Human.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P03372

### Technical Details

Immunogen	E.coli-derived human Estrogen Receptor recombinant protein (Position: F425-V595). Human Estrogen Receptor shares 89% and 88% amino acid (aa) sequences identity with mouse and rat Estrogen Receptor, respectively.
Predicted Reactive Species	Human
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, By Heat</p> <p>Western blot, 0.1-0.5ug/ml, Human</p>

## Anti-Estrogen Receptor/ESR1 Antibody Picoband™ (PB9192) Images




Figure 1. Western blot analysis of Estrogen Receptor using anti-Estrogen Receptor antibody (PB9192). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human MCF-7 whole cell lysates,

Lane 2: human T-47D whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Estrogen Receptor antigen affinity purified polyclonal antibody (Catalog # PB9192) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Estrogen Receptor at approximately 66 kDa. The expected band size for Estrogen Receptor is at 66 kDa.




Figure 2. IHC analysis of Estrogen Receptor using anti-Estrogen Receptor antibody (PB9192).

Estrogen Receptor was detected in a paraffin-embedded section of human mammary cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 ug/ml rabbit anti-Estrogen Receptor Antibody (PB9192) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.

## 5 Publications Citing This Product

1. PubMed ID: 25864222, Effects of estradiol on liver estrogen receptor-? and its mRNA expression in hepatic fibrosis in rats
2. PubMed ID: 29435117, An J, Liu T, She R, Wu Q, Tian J, Shi R, Hao W, Ren X, Yang Y, Lu Y, Yang Y, Wu Y. Oncotarget. 2017 Dec 17;9(4):4475-4484. doi: 10.18632/oncotarget.23381. eCollection 2018 Jan 12. Replication of hepatitis E virus in the ovary and promotion of oocy...
3. PubMed ID: 22101081, Fu Y, Zhao Z, Wu Y, Wu K, Xu X, Liu Y, Tong C. J Ethnopharmacol. 2012 Jan 6;139(1):26-33. Doi: 10.1016/J.Jep.2011.08.077. Epub 2011 Nov 10. Therapeutic Mechanisms Of Tongmai Dasheng Tablet On Tripterygium Glycosides Induced Rat Model For Premature...

Visit [bosterbio.com/anti-estrogen-receptor-picoband-trade-antibody-pb9192-boster.html](http://bosterbio.com/anti-estrogen-receptor-picoband-trade-antibody-pb9192-boster.html) to see all 5 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Estrogen Receptor/ESR1 Antibody <sup>™</sup>