

Anti-HOXA11 Antibody Picoband®

Catalog Number: PB9218

About HOXA11

Homeobox protein Hox-A11 is a protein that in humans is encoded by the HOXA11 gene. It is mapped to 7p15.2. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is involved in the regulation of uterine development and is required for female fertility. HOXA11 is essential for the development of the uterosacral ligaments, it has been found that women with pelvic organ prolapse may have weakened connective tissue due to changes in this signaling pathway. Mutations in this gene can cause radio-ulnar synostosis with amegakaryocytic thrombocytopenia.

Submit a product review to Biocompare.com

[Submit a review of this product](#) to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Overview

Product Name	Anti-HOXA11 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-HOXA11 Antibody Picoband® catalog # PB9218. Tested in Flow Cytometry, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	Flow Cytometry, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , and 0.05 mg NaN ₃ . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P31270


Technical Details

Immunogen	E.coli-derived human HOXA11 recombinant protein (Position: H74-R245). Human HOXA11 shares 95% amino acid (aa) sequence identity with mouse HOXA11.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(F) and ICC.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat Immunohistochemistry (Frozen Section), 0.5-1ug/ml, Human Immunocytochemistry, 0.5-1ug/ml, Human Flow Cytometry (Fixed), 1-3ug/1x10 ⁶ cells, Human


Anti-HOXA11 Antibody Picoband® (PB9218) Images



Western blot analysis of HOXA11 using anti-HOXA11 antibody (PB9218). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. lane 1: Recombinant Human HOXA11 Protein 0.5ng. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-HOXA11 antigen affinity purified polyclonal antibody (Catalog # PB9218) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for HOXA11 at approximately 38KD. The expected band size for HOXA11 is at 38KD.



Western blot analysis of HOXA11 using anti-HOXA11 antibody (PB9218). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: Rat Brain Tissue Lysate, Lane 2: Human Placenta Tissue Lysate, Lane 3: HLEA Whole Cell Lysate, Lane 4: HT1080 Whole Cell Lysate, Lane 5: HEPA Whole Cell Lysate. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-HOXA11 antigen affinity purified polyclonal antibody (Catalog # PB9218) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for HOXA11 at approximately 35KD. The expected band size for HOXA11 is at 35KD.



Flow Cytometry analysis of U937 cells using anti-HOXA11 antibody (PB9218). Overlay histogram showing U937 cells stained with PB9218 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-HOXA11 Antibody (PB9218, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight® 488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with

primary antibody and secondary antibody (Red line) was used as a blank control.

1 Publications Citing This Product

1. PubMed ID: 31699823, Xiong Z,Xia P,Zhu X,Geng J,Wang S,Ye B,Qin X,Qu Y,He L,Fan D,Du Y,Tian Y,Fan Z.Glutamylation of deubiquitinase BAP1 controls self-renewal of hematopoietic stem cells and hematopoiesis.J Exp Med.2020 Feb 3;217(2):e20190974.doi:10.1084/jem.20190974.PMID:31699823;PMCID:PMC7041701.

Visit bosterbio.com/anti-hoxa11-picoband-trade-antibody-pb9218-boster.html to see all 1 publications.

[Anti-HOXA11 Antibody](#)

For Research Use Only. Not for use in diagnostic procedures.