BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-HRG Antibody Picoband®

Catalog Number: PB9220

About HRG

Histidine-rich glycoprotein (HRG) is a protein that in humans is encoded by the HRG gene. It is mapped to 3q27.3. HRG contains two cystatin-like domains and is located in plasma and platelets. The protein can bind heme, dyes and divalent metal ions. It also can inhibit rosette formation and interacts with heparin, thrombospondin and plasminogen. Two of the protein's effects, the inhibition of fibrinolysis and the reduction of inhibition of coagulation, indicate a potential prothrombotic effect. Mutations in this gene lead to thrombophilia due to abnormal histidine-rich glycoprotein levels. HRG is thought to have multiple roles as a protein in the human blood, including roles in immunity, angiogenesis and coagulation.

Submit a product review to Biocompare.com



Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-HRG Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-HRG Antibody Picoband® catalog # PB9220. Tested in IHC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na2HPO4, and 0.05 mg NaN3. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20˚C for one year from date of receipt. After reconstitution, at 4˚C for one month. It can also be aliquotted and stored frozen at -20˚C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P04196


Technical Details

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Immunogen	E.coli-derived human HRG recombinant protein (Position: V19-V200). Human HRG shares 71% and 69% amino acid (aa) sequences identity with mouse and rat HRG, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human Western blot, 0.1-0.5ug/ml, Human

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


Anti-HRG Antibody Picoband® (PB9220) Images



Anti-HRG antibody, PB9220, Western blottingAll lanes: Anti HRG (PB9220) at 0.5ug/mlWB: Recombinant Human HRG Protein 0.5ngPredicted bind size: 40KDObserved bind size: 40KD



Anti-HRG antibody, PB9220, Western blottingAll lanes: Anti HRG (PB9220) at 0.5ug/mlWB: Human Placenta Tissue Lysate at 50ugPredicted bind size: 60KDObserved bind size: 60KD



Anti-HRG antibody, PB9220, IHC(P)IHC(P): Human Intestinal Cancer Tissue

Anti-HRG Antibody

For Research Use Only. Not for use in diagnostic procedures.