

Anti-Dystrophin/DMD Antibody Picoband® Biotin Conjugated

Catalog Number: PB9276-Biotin

About DMD

Dystrophin, also known as DMD, is a rod-shaped cytoplasmic protein, and a vital part of a protein complex that connects the cytoskeleton of a muscle fiber to the surrounding extracellular matrix through the cell membrane. It is mapped to Xp21.2-p21.1. This complex is variously known as the costamere or the dystrophin-associated protein complex. Many muscle proteins, such as alpha-dystrobrevin, syncoilin, synemin, sarcoglycan, dystroglycan, and sarcospan, colocalize with dystrophin at the costamere. Dystrophin is a protein located between the sarcolemma and the outermost layer of myofilaments in the muscle fiber (myofiber). It is a cohesive protein, linking actin filaments to another support protein that resides on the inside surface of each muscle fiber's plasma membrane (sarcolemma).

Overview

Product Name	Anti-Dystrophin/DMD Antibody Picoband® Biotin Conjugated
Reactive Species	Human, Mouse, Rat
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	P11532

Technical Details

Immunogen	E.coli-derived human Dystrophin recombinant protein (Position: H3076-D3404). Human Dystrophin shares 100% amino acid (aa) sequence identity with mouse Dystrophin.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	Biotin
Suggested Dilutions	The intended application should be selected according to the customer's experimental requirements.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Dystrophin/DMD Antibody - Biotin

For Research Use Only. Not for use in diagnostic procedures.