

Anti-Cyclin D1/CCND1 Antibody Picoband® FITC Conjugated

Catalog Number: PB9370-FITC

About CCND1

Cyclin D1, also known as CCND1, is a human gene. The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclin D1 encodes the regulatory subunit of a holoenzyme that phosphorylates and inactivates the retinoblastoma protein and promotes progression through the G1-S phase of the cell cycle. Amplification or overexpression of cyclin D1 plays pivotal roles in the development of a subset of human cancers including parathyroid adenoma, breast cancer, colon cancer, lymphoma, melanoma, and prostate cancer. The cyclin D1 gene is overexpressed in human breast cancers and is required for oncogene-induced tumorigenesis. Brisken et al. (2003) found that prolactin induced IGF2 mRNA and IGF2 induced cyclin D1 protein expression in mouse mammary epithelial cultures. And they also concluded that IGF2 is a mediator of prolactin-induced alveologenesis and that prolactin, IGF2, and cyclin D1 are components of a developmental pathway in mammary gland.

Overview

Product Name	Anti-Cyclin D1/CCND1 Antibody Picoband® FITC Conjugated
Reactive Species	Human, Mouse, Rat
Application	Flow Cytometry
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P24385

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human Cyclin D1, different from the related mouse sequence by two amino acids, and from the related rat sequence by one amino acid.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	FITC

	Excitation Wavelength: 495 nm Emission Wavelength: 525 nm
Suggested Dilutions	Flow Cytometry, Optimal dilutions should be determined by end users.

25 Publications Citing This Product

1. PubMed ID: 28944904, Triptolide inhibits proliferation, differentiation and induces apoptosis of osteoblastic MC3T3-E1 cells
2. PubMed ID: 27602489, Pivwil2 is reactivated by HPV oncoproteins and initiates cell reprogramming via epigenetic regulation during cervical cancer tumorigenesis
3. PubMed ID: 24788349, Zhang H, Zhao B, Huang C, Meng Xm, Bian Eb, Li J. Plos One. 2014 May 2;9(5):E95520. Doi: 10.1371/Journal.Pone.0095520. Ecollection 2014. Melittin Restores Pten Expression By Down-Regulating Hdac2 In Human Hepatocellular Carcinoma Hepg2 Cells.

Visit bosterbio.com/anti-cyclin-d1-picoband-trade-antibody-pb9370-boster.html to see all 25 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Cyclin D1/CCND1 Antibody - FITC

For Research Use Only. Not for use in diagnostic procedures.