

## Anti-CD36 Antibody Picoband®

Catalog Number: PB9371

### About CD36

CD36 (cluster of differentiation 36), also known as FAT (fatty acid translocase), FAT/CD36, (FAT)/CD36, SCARB3, GP88, glycoprotein IV (gpIV), and glycoprotein IIIb (gpIIIb), is an integral membrane protein found on the surface of many cell types in vertebrate animals. CD36 is a member of the class B scavenger receptor family of cell surface proteins. It is mapped to 7q21.11. And CD36 binds many ligands including collagen, thrombospondin, erythrocytes parasitized with Plasmodium falciparum, oxidized low density lipoprotein, native lipoproteins, oxidized phospholipids, and long-chain fatty acids. In addition, CD36 function in long-chain fatty acid uptake and signaling can be irreversibly inhibited by sulfo-N-succinimidyl oleate (SSO), which binds lysine 164 within a hydrophobic pocket shared by several CD36 ligands, e.g. fatty acid and oxLDL.

### Overview


Product Name	Anti-CD36 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-CD36 Antibody Picoband® catalog # PB9371. Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> , and 0.05 mg NaN <sub>3</sub> . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P16671

### Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human CD36, different from the related mouse sequence by six amino acids, and from the related rat sequence by four amino acids.
-----------	---

Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, Rat Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat


## Anti-CD36 Antibody Picoband® (PB9371) Images



Anti-CD36 Picoband antibody, PB9371, IHC(P)IHC(P): Rat Spleen Tissue



Anti-CD36 Picoband antibody, PB9371, Western blotting  
All lanes: Anti CD36 (PB9371) at 0.5ug/ml  
Lane 1: Rat Liver Tissue Lysate at 50ug  
Lane 2: Rat Cardiac Muscle Tissue Lysate at 50ug  
Lane 3: Mouse Liver Tissue Lysate at 50ug  
Lane 4: Mouse Cardiac Muscle Tissue Lysate at 50ug  
Lane 5: SMMC Whole Cell Lysate at 40ug  
Predicted bind size: 53KD  
Observed bind size: 88KD



IHC analysis of Galectin 1 using anti-Galectin 1 antibody (PB9371). Galectin 1 was detected in paraffin-embedded section of human Lung cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-Galectin 1 Antibody (PB9371) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

## 3 Publications Citing This Product

1. PubMed ID: 10.1159/000510032, Higher Levels of Triglyceride, Fatty Acid Translocase, and Toll-Like Receptor 4 and Lower Level of HDL-C in Pregnant Women with GDM and Their Close Correlation with Neonatal Weight
2. PubMed ID: 33819629, Huang Y,Zhao C,Kong Y,Tan P,Liu S,Liu Y,Zeng F,Yuan Y,Zhao B,Wang J.Elucidation of the mechanism of NEFA-induced PERK-eIF2alpha signaling pathway regulation of lipid metabolism in bovine hepatocytes.J Steroid Biochem Mol Biol.2021 Apr 2:105893.doi:10.1016/j.jsbmb.2021.105893.Epub ahead of print.PMID:33819629.
3. PubMed ID: 33205615, Kong Y, Zhao C, Huang Y, Liu Y, Liu S, Guo Y, Li M, Xu T, Zhao B, Wang J. Angiopoietin-like protein 4 promotes very-low-density lipoprotein assembly and secretion in bovine hepatocytes in vitro. IUBMB Life. 2020 Nov 17. doi: 10.1002/iub.2403. Epub ahead of

Visit [bosterbio.com/anti-cd36-picoband-trade-antibody-pb9371-boster.html](https://www.bosterbio.com/anti-cd36-picoband-trade-antibody-pb9371-boster.html) to see all 3 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-CD36 Antibody

For Research Use Only. Not for use in diagnostic procedures.