BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Synaptophysin/SYP Antibody Picoband®

Catalog Number: PB9409

About SYP

SYP (Synaptophysin) is located on the short arm of X chromosome (Xp11.23-p11.22). The protein is a synaptic vesicle glycoprotein with four transmembrane domains weighing 38kDa. It is present inneuroendocrine cells and in virtually all neurons in the brain and spinal cord that participate in synaptic transmission. It acts as a marker for neuroendocrine tumors, and its ubiquity at the synapse has led to the use of synaptophysin immunostaining for quantification of synapses. Also, the protein binds cholesterol and is thought to direct targeting of vesicle-associated membrane protein 2 (synaptobrevin) to intracellular compartments. Mutations in this gene are associated with X-linked mental retardation (XLMR).

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-Synaptophysin/SYP Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Synaptophysin/SYP Antibody Picoband® catalog # PB9409. Tested in WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na2HPO4, and 0.05 mg NaN3. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20˚C for one year from date of receipt. After reconstitution, at 4˚C for one month. It can also be aliquotted and stored frozen at -20˚C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P08247

BOSTER
antibody and ELISA experts


BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human Synaptophysin, identical to the related mouse and rat sequences.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Mouse, Rat, Human

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Synaptophysin/SYP Antibody Picoband® (PB9409) Images



Western blot analysis of Synaptophysin/SYP using anti-Synaptophysin/SYP antibody (PB9409). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat brain tissue lysates, Lane 2: rat hippocampus tissue lysates, Lane 3: mouse brain tissue lysates, Lane 4: mouse hippocampus tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Synaptophysin/SYP antigen affinity purified polyclonal antibody (PB9409) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054) at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for Synaptophysin/SYP at approximately 38 kDa. The expected band size for Synaptophysin/SYP is at 34 kDa.

3 Publications Citing This Product

1. PubMed ID: 24457043, Chen Y, Wei G, Nie H, Lin Y, Tian H, Liu Y, Yu X, Cheng S, Yan R, Wang Q, Liu Dh, Deng W, Lai Y, Zhou Jh, Zhang Sx, Lin Ww, Chen Df. Brain Res. 2014 Mar 13;1552:41-54. Doi: 10.1016/J.Brainres.2014.01.005. Epub 2014 Jan 20. ??-Asarone Prevents Auto...

2. PubMed ID: 25206583, Liu Y, Liang Z, Liu J, Zou W, Li X, Wang Y, An L. Neural Regen Res. 2013 Oct 15;8(29):2725-33. Doi: 10.3969/J.Issn.1673-5374.2013.29.004. Downregulation Of Caveolin-1 Contributes To The Synaptic Plasticity Deficit In The Hippocampus Of Aged Rats.

3. PubMed ID: 23271319, Li Xh, Xin X, Wang Y, Wu Jz, Jin Zd, Ma Ln, Nie Cj, Xiao X, Hu Y, Jin Mw. J Alzheimers Dis. 2013;34(3):755-67. Doi: 10.3233/Jad-122017. Pentamethylquercetin Protects Against Diabetes-Related Cognitive Deficits In Diabetic Goto-Kakizaki Rats.

Visit bosterbio.com/anti-synaptophysin-picoband-trade-antibody-pb9409-boster.html to see all 3 publications.

For Research Use Only. Not for use in diagnostic procedures.