

## Anti-SSH3BP1/ABI1 Antibody Picoband™

Catalog Number: PB9416

### About ABI1

SSH3BP1 is also known as ABI1. ABI1 is a human homolog of mouse Abl-interactor-1 (Abi1), mapped on 10p11.2. ABI1 participates in the transduction of signals from Ras to Rac by regulating Rac-specific guanine nucleotide exchange factor (GEF) activities. It dramatically promoted ABL1-mediated tyrosine phosphorylation of MENA, but not of other substrates. Abi-1 regulates c-Abl-mediated phosphorylation of Mena by interacting with both proteins. ABI1 plays a role in the leukemogenesis by translocating to MLL.

### Overview

Product Name	Anti-SSH3BP1/ABI1 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-SSH3BP1/ABI1 Antibody Picoband™ catalog # PB9416. Tested in Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IF, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q8IZP0

### Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human ABI1, identical to the related mouse and rat sequences.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and ICC.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat</p> <p>Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Mouse, Rat, By Heat</p> <p>Immunocytochemistry/Immunofluorescence, 5 ug/ml, Human</p> <p>Flow Cytometry, 1-3 ug/1x10<sup>6</sup> cells, Human</p>

## Anti-SSH3BP1/ABI1 Antibody Picoband™ (PB9416) Images




Figure 1. Western blot analysis of SSH3BP1/ABI1 using anti-SSH3BP1/ABI1 antibody (PB9416).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human 293T whole cell lysates,  
Lane 2: human MCF-7 whole cell lysates,  
Lane 3: human U87 whole cell lysates,  
Lane 4: rat brain tissue lysates,  
Lane 5: rat C6 whole cell lysates,  
Lane 6: mouse brain tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SSH3BP1/ABI1 antigen affinity purified polyclonal antibody (Catalog # PB9416) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for SSH3BP1/ABI1 at approximately 65 kDa. The expected band size for SSH3BP1/ABI1 is at 55 kDa.




Figure 2. IHC analysis of SSH3BP1/ABI1 using anti-SSH3BP1/ABI1 antibody (PB9416).

SSH3BP1/ABI1 was detected in a paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SSH3BP1/ABI1 Antibody (PB9416) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 3. IHC analysis of SSH3BP1/ABI1 using anti-SSH3BP1/ABI1 antibody (PB9416).

SSH3BP1/ABI1 was detected in a paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SSH3BP1/ABI1 Antibody (PB9416) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog #

SV0002) with DAB as the chromogen.




Figure 4. IF analysis of SSH3BP1/ABI1 using anti-SSH3BP1/ABI1 antibody (PB9416). SSH3BP1/ABI1 was detected in an immunocytochemical section of A431 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-SSH3BP1/ABI1 Antibody (PB9416) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 5. Flow Cytometry analysis of K562 cells using anti-SSH3BP1/ABI1 antibody (PB9416). Overlay histogram showing K562 cells stained with PB9416 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-SSH3BP1/ABI1 Antibody (PB9416, 1 ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

## 1 Publications Citing This Product

1. PubMed ID: 19554484, Cui M, Yu W, Dong J, Chen J, Zhang X, Liu Y. Med Oncol. 2010 Sep;27(3):632-9. Doi: 10.1007/S12032-009-9260-6. Epub 2009 Jun 25. Downregulation Of Abi1 Expression Affects The Progression And Prognosis Of Human Gastric Carcinoma.

Visit [bosterbio.com/anti-abi1-picoband-trade-antibody-pb9416-boster.html](http://bosterbio.com/anti-abi1-picoband-trade-antibody-pb9416-boster.html) to see all 1 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-SSH3BP1/ABI1 Antibody <sup>™</sup>