

Anti-Aryl hydrocarbon Receptor/AHR Antibody Picoband® FITC Conjugated

Catalog Number: PB9419-FITC

About Ahr

AHR (aryl hydrocarbon receptor), also called bHLHe76, is a member of the family of basic helix-loop-helix transcription factors. AhR is a cytosolic transcription factor that is normally inactive, bound to several co-chaperones. The AHR gene is mapped on 7p21.1. Estrogenic actions of AHR agonists were detected in wildtype ovariectomized mouse uteri, but were absent in Ahr -/- or Er-alpha -/- ovariectomized mice. Complex assembly and ubiquitin ligase activity of CUL4B (AHR) in vitro and in vivo are dependent on the AHR ligand. In the CUL4B (AHR) complex, ligand-activated AHR acts as a substrate-specific adaptor component that targets sex steroid receptors for degradation. Cd4-positive cells from mice lacking Ahr developed Th17 responses but failed to produce IL22 and did not show enhanced Th17 development. Activation of Ahr during induction of EAE accelerated disease onset and increased pathology in wildtype mice, but not in Ahr -/- mice. The TDO-AHR pathway is active in human brain tumors and is associated with malignant progression and poor survival. Ahr activity within ROR-gamma-t-positive ILC could be induced by dietary ligands such as those contained in vegetables of the family Brassicaceae.

Overview

Product Name	Anti-Aryl hydrocarbon Receptor/AHR Antibody Picoband® FITC Conjugated
Reactive Species	Mouse, Rat
Application	Recommended applications are based on the parent unconjugated antibody (WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na2HPO4, 0.02% NaN3.
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P30561

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of mouse AHR, different from the related human sequence by five amino acids, and from the related rat sequence by one amino acid.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL

Purification	Immunogen affinity purified.
Conjugate	FITC Excitation Wavelength: 495 nm Emission Wavelength: 525 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

1 Publications Citing This Product

1. PubMed ID: 33204326, Yang X,Liu H,Ye T,Duan C,Lv P,Wu X,Liu J,Jiang K,Lu H,Yang H,Xia D,Peng E,Chen Z,Tang K,Ye Z. AhR activation attenuates calcium oxalate nephrocalcinosis by diminishing M1 macrophage polarization and promoting M2 macrophage polarization. Theranostics.2020

Visit bosterbio.com/anti-ahr-picoband-trade-antibody-pb9419-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Aryl hydrocarbon Receptor/AHR Antibody - FITC

For Research Use Only. Not for use in diagnostic procedures.