

Anti-SNAI3 Antibody Picoband®

Catalog Number: PB9440

About SNAI3

SNAI3 is a member of the SNAIL gene family, named for the *Drosophila* snail gene, which plays roles in mesodermal formation during embryogenesis. SNAI3 gene, was located between KIAA0233 gene and CBFA2T3 gene in human chromosome 16q24.3, a region affected in breast cancer, gastric cancer, hepatocellular carcinoma, ovarian cancer, and therapy-related myeloid leukemia with t(16;21)(q24;q22) translocation. Human SNAI3 gene was found to encode 292-amino-acid polypeptide with the N-terminal SNAG domain and five zinc finger domains. And Human SNAI3 mRNA was expressed in skin melanotic melanoma, lung epidermoid carcinoma, and germ cell tumor. Because SNAG zinc-finger proteins are transcriptional repressors implicated in carcinogenesis and embryogenesis, SNAI3 gene might be a potent target of pharmacogenomics in the field of oncology and regenerative medicine.

Overview

Product Name	Anti-SNAI3 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-SNAI3 Antibody Picoband® catalog # PB9440. Tested in WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , and 0.05 mg NaN ₃ . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q3KNW1

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human SNAI3, different from the related mouse sequence by two amino acids.
-----------	--

Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human

Anti-SNAI3 Antibody Picoband® (PB9440) Images



Western blot analysis of SNAI3 using anti-SNAI3 antibody (PB9440). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. Lane 1: MCF-7 Whole Cell Lysate at 40ug. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SNAI3 antigen affinity purified polyclonal antibody (Catalog # PB9440) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for SNAI3 at approximately 32 kDa. The expected band size for SNAI3 is at 32 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-SNAI3 Antibody

For Research Use Only. Not for use in diagnostic procedures.