BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Cystathionase/CTH Antibody Picoband®

Catalog Number: PB9494

About CTH

Cystathionine gamma-lyase (or cystathionase) is an enzyme which breaks down cystathionine into cysteine and alpha-ketobutyrate. The International Radiation Hybrid Mapping Consortium mapped the CTH gene to chromosome 1. The CTH gene had earlier been assigned to chromosome 16 by study of somatic cell hybrids. It is demonstrated that hydrogen sulfide (H2S) is physiologically generated by CTH.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-Cystathionase/CTH Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Cystathionase/CTH Antibody Picoband® catalog # PB9494. Tested in WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na2HPO4, and 0.05 mg NaN3. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20˚C for one year from date of receipt. After reconstitution, at 4˚C for one month. It can also be aliquotted and stored frozen at -20˚C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P32929


Technical Details

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com


Immunogen	E.coli-derived human Cystathionase recombinant protein (Position: D181-H398). Human Cystathionase shares 85.8% amino acid (aa) sequence identity with both mouse and rat Cystathionase.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-Cystathionase/CTH Antibody Picoband® (PB9494) Images



Western blot analysis of Cystathionase using anti-Cystathionase antibody (PB9494). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. lane 1: K562 whole cell lysates, lane 2: HepG2 whole cell lysates, lane 3: RH35 whole cell lysates, lane 4: rat liver tissue lysates, lane 5: rat kidney tissue lysates, lane 6: mouse liver tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Cystathionase antigen affinity purified polyclonal antibody (Catalog # PB9494) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Cystathionase at approximately 45KD. The expected band size for Cystathionase is at 45KD.



Western blot analysis of Gamma Cystathionase/CTH using anti-Gamma Cystathionase/CTH antibody (PB9494). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2- WT whole cell lysates, Lane 2: human HepG2-CTH KO whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Gamma Cystathionase/CTH antigen affinity purified polyclonal antibody (PB9494) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for Gamma Cystathionase/CTH at approximately 42 kDa. The expected band size for Gamma Cystathionase/CTH is at 45 kDa.

6 Publications Citing This Product

1. PubMed ID: 33649809, Nie L,Liu M,Chen J,Wu Q,Li Y,Yi J,Zheng X,Zhang J,Chu C,Yang J.Hydrogen sulfide ameliorates doxorubicin-induced myocardial fibrosis in rats via the PI3K/AKT/mTOR pathway.Mol Med Rep.2021 Apr;23(4):299.doi:10.3892/mmr.2021.11938.Epub 2021 Mar 2.PMID:336498

2. PubMed ID: 29393353, Hydrogen sulfide attenuates myocardial fibrosis in diabetic rats through the JAK/STAT signaling pathway

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

3. PubMed ID: 26378818, Caffeic Acid Phenethyl Ester inhibit Hepatic Fibrosis by Nitric Oxide Synthase and Cystathionine Gamma-Lyase in Rats

Visit bosterbio.com/anti-cystathionase-picoband-trade-antibody-pb9494-boster.html to see all 6 publications.

Anti-Cystathionase/CTH Antibody

For Research Use Only. Not for use in diagnostic procedures.