

Anti-Transcription factor Sp4 Antibody Picoband®

Catalog Number: PB9558

About SP4

SP4 (Sp4 transcription factor), also known as Transcription factor Sp4, MGC130008, MGC130009, SPR-1, HF1B, is a protein that in humans is encoded by the SP4 gene. It is a multigene family of SP proteins. Northern blot analysis that SP4 is abundantly expressed as a 7.5-kb message in brain, but barely detectable in other human cell lines. SP4 is an activator protein like SP1. However, in contrast to SP1, SP4 is not able to act synergistically through adjacent binding sites.

Overview


Product Name	Anti-Transcription factor Sp4 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Transcription factor Sp4 Antibody Picoband® catalog # PB9558. Tested in WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q02446

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human SP4, different from the related mouse sequence by one amino acid.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat

Anti-Transcription factor Sp4 Antibody Picoband® (PB9558) Images



Western blot analysis of SP4 using anti-SP4 antibody (PB9558). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. Lane 1: Rat Brain Tissue Lysate at 50ug, Lane 2: Mouse Brain Tissue Lysate at 50ug, Lane 3: U87 Whole Cell Lysate at 40ug, Lane 4: HELA Whole Cell Lysate at 40ug. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SP4 antigen affinity purified polyclonal antibody (Catalog # PB9558) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for SP4 at approximately 82 kDa. The expected band size for SP4 is at 82 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Transcription factor Sp4 Antibody

For Research Use Only. Not for use in diagnostic procedures.