

Anti-Eph receptor A5/EPHA5 Antibody Picoband™

Catalog Number: PB9583

About EPHA5

EPH receptor A5 (ephrin type-A receptor 5) is a protein that in humans is encoded by the EPHA5 gene. This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Alternatively spliced transcript variants encoding different isoforms have been described.

Overview

Product Name	Anti-Eph receptor A5/EPHA5 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Eph receptor A5/EPHA5 Antibody Picoband™ catalog # PB9583. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P54756

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Eph receptor A5, different from the related mouse sequence by two amino acids.
Predicted Reactive Species	Chicken
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Mouse, Rat, Human</p>

Anti-Eph receptor A5/EPHA5 Antibody Picoband™ (PB9583) Images




Figure 1. Western blot analysis of Eph receptor A5 using anti-Eph receptor A5 antibody (PB9583).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours.

Lane 1: Rat Brain Tissue Lysate at 50ug

Lane 2: Mouse Brain Tissue Lysate at 50ug.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5

hour at RT. The membrane was incubated with rabbit anti-

Eph receptor A5 antigen affinity purified polyclonal antibody

(Catalog # PB9583) at 0.5 ug/mL overnight at 4°C, then

washed with TBS-0.1%Tween 3 times with 5 minutes each

and probed with a goat anti-rabbit IgG-HRP secondary

antibody at a dilution of 1:5000 for 1.5 hour at RT. The

signal is developed using an Enhanced Chemiluminescent

detection (ECL) kit (Catalog # EK1002) with Tanon 5200

system. A specific band was detected for Eph receptor A5 at

approximately 114 kDa. The expected band size for Eph

receptor A5 is at 114 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Eph receptor A5/EPHA5 Antibody [™]