

Anti-Fascin/FSCN1 Antibody Picoband®

Catalog Number: PB9592

About FSCN1

Fascin is an actin cross-linking protein. The Fascin gene contains 5 exons and spans 7 kb. It is a 54-58 kilodalton monomeric actin filament bundling protein originally isolated from sea urchin egg but also found in Drosophila and vertebrates, including humans. Fascin (from the Latin for bundle) is spaced at 11 nanometre intervals along the filament. The bundles in cross section are seen to be hexagonally packed, and the longitudinal spacing is compatible with a model where fascin cross-links at alternating 4 and 5 actins. It is calcium insensitive and monomeric. Fascin binds beta-catenin, and colocalizes with it at the leading edges and borders of epithelial and endothelial cells. The role of Fascin in regulating cytoskeletal structures for the maintenance of cell adhesion, coordinating motility and invasion through interactions with signalling pathways is an active area of research especially from the cancer biology perspective. Abnormal fascin expression or function has been implicated in breast cancer, colon cancer, esophageal squamous cell carcinoma, gallbladder cancer and prostate cancer.

Overview

Product Name	Anti-Fascin/FSCN1 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Fascin/FSCN1 Antibody Picoband® catalog # PB9592. Tested in WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl and 0.2mg Na2HPO4.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q16658

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human Fascin, identical to the related mouse sequence, and different from the related rat sequence by one amino acid.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins





BOSTER
antibody and ELISA experts

Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat



Anti-Fascin/FSCN1 Antibody Picoband® (PB9592) Images




Figure 1. Western blot analysis of Fascin/FSCN1 using anti-Fascin/FSCN1 antibody (PB9592).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human SH-SY5Y whole cell lysates,

Lane 2: human Hela whole cell lysates,

Lane 3: human K562 whole cell lysates,

Lane 4: human A549 whole cell lysates,

Lane 5: human HepG2 whole cell lysates,

Lane 6: human PC-3 whole cell lysates,

Lane 7: rat brain tissue lysates,

Lane 8: mouse brain tissue lysates,

Lane 9: mouse NIH/3T3 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Fascin/FSCN1 antigen affinity purified polyclonal antibody (Catalog # PB9592) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Fascin/FSCN1 at approximately 55 kDa. The expected band size for Fascin/FSCN1 is at 55 kDa.

1 Publications Citing This Product

1. PubMed ID: 25231007, Jiang L, Wang P, Chen H. Ups J Med Sci. 2014 Nov;119(4):324-32. Doi: 10.3109/03009734.2014.960053. Epub 2014 Sep 18. Overexpression Of Foxm1 Is Associated With Metastases Of Nasopharyngeal Carcinoma.

Visit bosterbio.com/anti-fascin-picoband-trade-antibody-pb9592-boster.html to see all 1 publications.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Fascin/FSCN1 Antibody