

## Anti-GSTA1/GSTA2/GSTA3/GSTA4/GSTA5 Antibody Picoband® Fluoro550 Conjugated

Catalog Number: PB9627-Fluoro550

### About GSTA1

Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. These enzymes function in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione. The genes encoding these enzymes are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of some drugs. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. This gene encodes a glutathione S-transferase belonging to the alpha class. The alpha class genes, located in a cluster mapped to chromosome 6, are the most abundantly expressed glutathione S-transferases in liver (hepatocytes) and kidney (proximal tubules). In addition to metabolizing bilirubin and certain anti-cancer drugs in the liver, the alpha class of these enzymes exhibit glutathione peroxidase activity, thereby protecting the cells from reactive oxygen species and the products of peroxidation.

### Overview

Product Name	Anti-GSTA1/GSTA2/GSTA3/GSTA4/GSTA5 Antibody Picoband® Fluoro550 Conjugated
Reactive Species	Human, Mouse, Rat
Application	Flow Cytometry
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na <sub>2</sub> HPO <sub>4</sub> , 0.02% NaN <sub>3</sub> .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P08263

### Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human GSTA1/A2/A3/A4/A5, different from the related mouse and rat sequences by five amino acids.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.

Conjugate	Fluoro550 Excitation Wavelength: 562 nm Emission Wavelength: 576 nm
Suggested Dilutions	Flow Cytometry, Optimal dilutions should be determined by end users.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-GSTA1/GSTA2/GSTA3/GSTA4/GSTA5 Antibody - Fluoro550

For Research Use Only. Not for use in diagnostic procedures.