

Anti-BMP4 Antibody Picoband®

Catalog Number: PB9688

About BMP4

Bone morphogenetic protein 4 is a protein that in humans is encoded by BMP4 gene. It is found on chromosome 14q22-q23. The protein encoded by this gene is a member of the bone morphogenetic protein family which is part of the transforming growth factor-beta superfamily. The superfamily includes large families of growth and differentiation factors. Bone morphogenetic proteins were originally identified by an ability of demineralized bone extract to induce endochondral osteogenesis in vivo in an extraskelatal site. This particular family member plays an important role in the onset of endochondral bone formation in humans, and a reduction in expression has been associated with a variety of bone diseases, including the heritable disorder Fibrodysplasia Ossificans Progressiva. Alternative splicing in the 5' untranslated region of this gene has been described and three variants are described, all encoding an identical protein.

Overview


Product Name	Anti-BMP4 Antibody Picoband®
Reactive Species	Human, Mouse
Description	Boster Bio Anti-BMP4 Antibody Picoband® catalog # PB9688. Tested in ELISA, WB applications. This antibody reacts with Human, Mouse. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , and 0.05 mg NaN ₃ . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P12644

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human BMP-4, different from the related mouse and rat sequences by two amino acids.
-----------	--

Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	ELISA, 0.1-0.5ug/ml, -, Mouse, - Western blot, 0.1-0.5ug/ml, Human, Mouse

Anti-BMP4 Antibody Picoband® (PB9688) Images



Western blot analysis of BMP-4 using anti-BMP-4 antibody (PB9688). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. Lane 1: Mouse Lung Tissue Lysate at 50ug, Lane 2: Mouse Liver Tissue Lysate at 50ug, Lane 3: HEPA Whole Cell Lysate at 40ug, Lane 4: HEPG2 Whole Cell Lysate at 40ug, Lane 5: HELA Whole Cell Lysate at 40ug. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-BMP-4 antigen affinity purified polyclonal antibody (Catalog # PB9688) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for BMP-4 at approximately 46 kDa. The expected band size for BMP-4 is at 46 kDa.

2 Publications Citing This Product

1. PubMed ID: 25709905, Wang Q, Xue MI, Zhao Gq, Liu Mg, Ma Yn, Ma Y. Int J Ophthalmol. 2015 Feb 18;8(1):39-45. Doi: 10.3980/J.Issn.2222-3959.2015.01.07. Ecollection 2015. Form-Deprivation Myopia Induces Decreased Expression Of Bone Morphogenetic Protein-2, 5 In Guinea P...
2. PubMed ID: 16336834, Fu Jr, Liu WI, Zhou Yf, Zhou Jf, Sun Hy, Luo L, Zhang H, Xu Hz. Chin Med J (Engl). 2005 Dec 5;118(23):1979-86. Expansive Effects Of Aorta-Gonad-Mesonephros-Derived Stromal Cells On Hematopoietic Stem Cells From Embryonic Stem Cells.

Visit bosterbio.com/anti-bmp-4-picoband-trade-antibody-pb9688-boster.html to see all 2 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-BMP4 Antibody

For Research Use Only. Not for use in diagnostic procedures.