

Anti-BCA1/CXCL13 Antibody Picoband®

Catalog Number: PB9697

About CXCL13

C-X-C motif chemokine 13 (CXCL13), also known as B lymphocyte chemoattractant (BLC), is a protein that in humans is encoded by the CXCL13 gene. CXCL13 is a small cytokine belonging to the CXC chemokine family. Its gene is mapped to human chromosome 4 in a cluster of other CXC chemokines. This protein has contained 109 amino acids. It is expressed highly in the liver, spleen, lymph nodes and gut of humans. The protein induces chemotaxis of BLR1-expressing B cells.

Overview

Product Name	Anti-BCA1/CXCL13 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-BCA1/CXCL13 Antibody Picoband® catalog # PB9697. Tested in WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O43927

Technical Details

Immunogen	E. coli-derived human BCA1 recombinant protein (Position: V23-P109). Human BCA1 shares 45.3% amino acid (aa) sequence identity with mouse BCA1.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human

Anti-BCA1/CXCL13 Antibody Picoband® (PB9697) Images



Western blot analysis of CXCL13 using anti-CXCL13 antibody (PB9697). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane : HELA whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CXCL13 antigen affinity purified polyclonal antibody (Catalog # PB9697) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CXCL13 at approximately 13KD. The expected band size for CXCL13 is at 13KD.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-BCA1/CXCL13 Antibody

For Research Use Only. Not for use in diagnostic procedures.