

Anti-FMO2 Antibody Picoband®

Catalog Number: PB9760

About FMO2

Dimethylaniline monooxygenase [N-oxide-forming] 2 is an enzyme that in humans is encoded by the FMO2 gene. This gene encodes a flavin-containing monooxygenase family member. It is an NADPH-dependent enzyme that catalyzes the N-oxidation of some primary alkylamines through an N-hydroxylamine intermediate. However, some human populations contain an allele (FMO2*2A) with a premature stop codon, resulting in a protein that is C-terminally-truncated, has no catalytic activity, and is likely degraded rapidly. This gene is found in a cluster with other related family members on chromosome 1. Alternative splicing results in multiple transcript variants.

Overview

Product Name	Anti-FMO2 Antibody Picoband®
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-FMO2 Antibody Picoband® catalog # PB9760. Tested in WB applications. This antibody reacts with Human;Mouse;Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q99518


Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human FMO2, different from the related mouse and rat sequences by two amino acids.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.

Suggested Dilutions

Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat

Anti-FMO2 Antibody Picoband® (PB9760) Images



Western blot analysis of FMO2 using anti-FMO2 antibody (PB9760). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat lung tissue lysates, Lane 2: mouse lung tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-FMO2 antigen affinity purified polyclonal antibody (PB9760) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054) at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for FMO2 at approximately 61 kDa. The expected band size for FMO2 is at 61 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-FMO2 Antibody

For Research Use Only. Not for use in diagnostic procedures.