

Anti-PI 3 Kinase p85 beta/PIK3R2 Antibody Picoband®

Catalog Number: PB9777

About PIK3R2

PIK3R2 (Phosphatidylinositol 3-kinase, regulatory subunit 2), also called p85-Beta, is an enzyme that in humans is encoded by the PIK3R2 gene. The PIK3R2 gene is mapped on 19p13.11. Phosphatidylinositol 3-kinase (PI3K) is a lipid kinase that phosphorylates phosphatidylinositol and similar compounds, creating second messengers important in growth signaling pathways. PI3K functions as a heterodimer of a regulatory and a catalytic subunit. The protein encoded by this gene is a regulatory component of PI3K. Two transcript variants, one protein coding and the other non-protein coding, have been found for this gene.

Overview


Product Name	Anti-PI 3 Kinase p85 beta/PIK3R2 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-PI 3 Kinase p85 beta/PIK3R2 Antibody Picoband® catalog # PB9777. Tested in WB, Flow Cytometry applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O00459

Technical Details


Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human PIK3R2, different from the related mouse sequence by two amino acids, and from the related rat sequence by one amino acid.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat Flow Cytometry(Fixed), 1-3 ug/1x10 ⁶ cells, Human

Anti-PI 3 Kinase p85 beta/PIK3R2 Antibody Picoband® (PB9777) Images



Western blot analysis of PIK3R2 using anti-PIK3R2 antibody (PB9777). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human A549 whole cell lysates, Lane 2: human Jurkat whole cell lysates, Lane 3: human K562 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PIK3R2 antigen affinity purified polyclonal antibody (Catalog # PB9777) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for PIK3R2 at approximately 82 kDa. The expected band size for PIK3R2 is at 82 kDa.



Flow Cytometry analysis of THP-1 cells using anti-PIK3R2 antibody (PB9777). Overlay histogram showing THP-1 cells stained with PB9777 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-PIK3R2 Antibody (PB9777, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-PI 3 Kinase p85 beta/PIK3R2 Antibody

For Research Use Only. Not for use in diagnostic procedures.