

## Anti-TRAP1 Antibody Picoband®

Catalog Number: PB9834

### About TRAP1

Heat shock protein 75 kDa, mitochondrial is a protein that in humans is encoded by the TRAP1 gene. It is mapped to 16p13.3. This gene encodes a mitochondrial chaperone protein that is member of the heat shock protein 90 (HSP90) family. The encoded protein has ATPase activity and interacts with tumor necrosis factor type I. And this protein may function in regulating cellular stress responses. In addition, it was found that TRAP1 interacted with the N-terminal half of TNFR1. Also, TRAP1 interacted with the C-terminal ends of the proteins encoded by both multiple exostoses-causing genes, EXT1 and EXT2, but not with EXTL1 or EXTL3.

### Overview


Product Name	Anti-TRAP1 Antibody Picoband®
Reactive Species	Human, Rat
Description	Boster Bio Anti-TRAP1 Antibody Picoband® catalog # PB9834. Tested in IHC, WB applications. This antibody reacts with Human, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains antibody formulated with stabilizing components, 0.9 mg NaCl, 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> , and 0.05 mg NaN <sub>3</sub> . *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q12931

### Technical Details


Immunogen	E.coli-derived human TRAP1 recombinant protein (Position: A571-H704). Human TRAP1 shares 91.7% and 94% amino acid (aa) sequence identity with mouse and rat TRAP1, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).

Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human Western blot, 0.1-0.5ug/ml, Human, Rat

## Anti-TRAP1 Antibody Picoband® (PB9834) Images



Western blot analysis of TRAP1 using anti-TRAP1 antibody (PB9834). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. Lane 1: Rat Cardiac Muscle Tissue Lysate at 50ug, Lane 2: Rat Kidney Tissue Lysate at 50ug, Lane 3: Rat Brain Tissue Lysate at 50ug, Lane 4: SMMC Whole Cell Lysate at 40ug, Lane 5: PANC Whole Cell Lysate at 40ug, Lane 6: A549 Whole Cell Lysate at 40ug. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TRAP1 antigen affinity purified polyclonal antibody (Catalog # PB9834) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for TRAP1 at approximately 80 kDa. The expected band size for TRAP1 is at 80 kDa.



IHC analysis of TRAP1 using anti-TRAP1 antibody (PB9834). TRAP1 was detected in a paraffin-embedded section of human intestinal cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 ug/ml rabbit anti-TRAP1 Antibody (PB9834) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TRAP1 Antibody

For Research Use Only. Not for use in diagnostic procedures.