

## Anti-CD33 Antibody Picoband®

Catalog Number: PB9932

### About CD33

CD33, also known as Siglec-3 (sialic acid binding Ig-like lectin 3, SIGLEC3, SIGLEC-3, gp67, p67), is a transmembrane receptor expressed on cells of myeloid lineage. It is usually considered myeloid-specific, but it can also be found on some lymphoid cells. CD33 binds sialic acids, therefore is a member of the SIGLEC family of lectins. By fluorescence in situ hybridization, CD33 is mapped to 19q13.3-q13.4.

### Overview


|                      |  |
|----------------------|--|
| Product Name         | Anti-CD33 Antibody Picoband®   |
| Reactive Species     | Human  |
| Description          | Boster Bio Anti-CD33 Antibody Picoband® catalog # PB9932. Tested in IHC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance. |
| Application          | IHC, WB  |
| Clonality            | Polyclonal   |
| Formulation          | Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .   |
| Storage Instructions | Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.  |
| Host                 | Rabbit   |
| Uniprot ID           | P20138   |

### Technical Details


|                               |  |
|-------------------------------|--|
| Immunogen                     | E. coli-derived human CD33 recombinant protein (Position: D18-H259). Human CD33 shares 61.6% amino acid (aa) sequence identity with mouse CD33.  |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P), IHC(F) and ICC. |
| Cross Reactivity              | No cross-reactivity with other proteins  |
| Isotype                       | Rabbit IgG   |
| Form                          | Lyophilized  |
| Concentration                 | Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.  |

|                     |  |
|---------------------|--|
| Purification        | Immunogen affinity purified.   |
| Suggested Dilutions | Western blot, 0.1-0.5ug/ml, Human<br>Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Human |

## Anti-CD33 Antibody Picoband® (PB9932) Images



Western blot analysis of CD33 using anti-CD33 antibody (PB9932). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human THP-1 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CD33 antigen affinity purified polyclonal antibody (PB9932) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054) at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for CD33 at approximately 60-75 kDa. The expected band size for CD33 is at 40 kDa.



IHC analysis of CD33 using anti-CD33 antibody (PB9932). CD33 was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-CD33 Antibody (PB9932) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

### Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-CD33 Antibody

For Research Use Only. Not for use in diagnostic procedures.