

Anti-Alpha A Crystallin/CRYAA Antibody Picoband® APC Conjugated

Catalog Number: PB9940-APC

About CRYAA

Alpha-crystallin A chain is a protein that in humans is encoded by the CRYAA gene. Mammalian lens crystallins are divided into alpha, beta, and gamma families. Alpha crystallins are composed of two gene products: alpha-A and alpha-B, for acidic and basic, respectively. Alpha crystallins can be induced by heat shock and are members of the small heat shock protein (HSP20) family. They act as molecular chaperones although they do not renature proteins and release them in the fashion of a true chaperone; instead they hold them in large soluble aggregates. Two additional functions of alpha crystallins are an autokinase activity and participation in the intracellular architecture. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Alpha-A and alpha-B gene products are differentially expressed; alpha-A is preferentially restricted to the lens and alpha-B is expressed widely in many tissues and organs. Defects in this gene cause autosomal dominant congenital cataract (ADCC).

Overview

Product Name	Anti-Alpha A Crystallin/CRYAA Antibody Picoband® APC Conjugated
Reactive Species	Human, Mouse, Rat
Application	Recommended applications are based on the parent unconjugated antibody (Flow Cytometry, IHC, IHC-F, ICC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P02489

Technical Details

Immunogen	E. coli-derived human Alpha A Crystallin recombinant protein (Position: M1-S173). Human Alpha A Crystallin shares 94.8% amino acid (aa) sequence identity with both mouse and rat Alpha A Crystallin.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.

Conjugate	APC Excitation Wavelength: 633-647 nm Emission Wavelength: 660 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Alpha A Crystallin/CRYAA Antibody - APC

For Research Use Only. Not for use in diagnostic procedures.