

Anti-GIP Antibody Picoband® Biotin Conjugated

Catalog Number: PB9947-Biotin

About Gip

Gastric inhibitory polypeptide (GIP), also known as the glucose-dependent insulinotropic peptide, is an inhibiting hormone of the secretin family of hormones. GIP is thought to have significant effects on fatty acid metabolism through stimulation of lipoprotein lipase activity in adipocytes. Additionally, GIP release has been demonstrated in the ruminant animal and may play a role in nutrient partitioning in milk production (lipid metabolism). Recently, GIP appeared as a major player in bone remodelling. It was evidenced that genetic ablation of the GIP receptor in mice resulted in profound alterations of bone microarchitecture through modification of the adipokine network. Furthermore, the deficiency in GIP receptors has also been associated in mice with a dramatic decrease in bone quality and a subsequent increase in fracture risk.

Overview

Product Name	Anti-GIP Antibody Picoband® Biotin Conjugated
Reactive Species	Mouse, Rat
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	P48756

Technical Details

Immunogen	E. coli-derived mouse GIP recombinant protein (Position: Y44-Q85). Mouse GIP shares 92.9% amino acid (aa) sequence identity with both human and rat GIP.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	Biotin
Suggested Dilutions	The intended application should be selected according to the customer's experimental requirements.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-GIP Antibody - Biotin

For Research Use Only. Not for use in diagnostic procedures.