

Anti-BCA1/CXCL13 Antibody Picoband™

Catalog Number: PB9999

About Cxcl13

C-X-C motif chemokine 13 (CXCL13), also known as B cell-attracting chemokine 1 (BCA-1), is a protein that in humans is encoded by the CXCL13 gene. CXCL13 is a small cytokine belonging to the CXC chemokine family. Its gene is mapped to human chromosome 4 in a cluster of other CXC chemokines. This protein has contained 109 amino acids. It is expressed highly in the liver, spleen, lymph nodes and gut of humans. The protein induces chemotaxis of BLR1-expressing B cells.

Overview

Product Name	Anti-BCA1/CXCL13 Antibody Picoband™
Reactive Species	Mouse
Description	Boster Bio Anti-BCA1/CXCL13 Antibody Picoband™ catalog # PB9999. Tested in ELISA, IHC, WB applications. This antibody reacts with Mouse.
Application	ELISA, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2mg Na2HPO4, 0.05 mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O55038

Technical Details

Immunogen	E. coli-derived mouse BCA1 recombinant protein (Position: I22-A109). Mouse BCA1 shares 45.3% amino acid (aa) sequence identity with human BCA1.
Predicted Reactive Species	Bovine
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.



BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Mouse, By Heat ELISA, 0.1-0.5ug/ml, Mouse, - Western blot, 0.1-0.5ug/ml, Mouse



Anti-BCA1/CXCL13 Antibody Picoband™ (PB9999) Images

70KD — 55KD —

100KD -

35KD-

15KD - _

Figure 1. Western blot analysis of CXCL13/BLC using anti-CXCL13/BLC antibody (PB9999).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. Lane 1: Recombinant mouse CXCL13/BLC protein 1ng.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CXCL13/BLC antigen affinity purified polyclonal antibody (Catalog # PB9999) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CXCL13/BLC at approximately 13 kDa. The expected band size for CXCL13/BLC is at 12 kDa.




Figure 2. IHC analysis of CXCL13/BLC using anti-CXCL13/BLC antibody (PB9999).

CXCL13/BLC was detected in a paraffin-embedded section of mouse lymphaden tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 ug/ml rabbit anti-CXCL13/BLC Antibody (PB9999) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-BCA1/CXCL13 Antibody ™