

Human recombinant FGF-2 (Fibroblast Growth Factors 2) protein, AF, Tag free

Catalog Number: PROTP09038-10

Overview

Product Name	Human recombinant FGF-2 (Fibroblast Growth Factors 2) protein, AF, Tag free
Description	Basic fibroblast Growth Factors (FGF-2, bFGF, FGF-beta), a 18 kDa pleiotropic cytokine, plays multiple roles in different cells and tissues. FGF-2 can stimulate smooth muscle cell growth, wound healing, and tissue repair. In addition, FGF-2 has been shown to regulate the generation of neurons and astrocytes from progenitor cells. FGF-2 are also involved in a variety of biological processes, including embryonic development, morphogenesis, tissue repair, tumor growth, and invasion. As a multifunctional cytokine, FGF-2 is first isolated from the pituitary. Later, it was identified from various cell types including cardiac myocytes, cardiac fibroblasts, endothelial cells, and smooth muscle cells.
Size	20ug,100ug,500ug
Tag	Tag Free
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.

Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Purity

>98% as determined by SDS-PAGE.

Molecular weight

The protein has a calculated MW of 18.1 kDa. The protein migrates as 15-18 kDa under reducing condition (SDS-PAGE analysis).

Activity

Measure by its ability to induce 3T3 cells proliferation. The ED₅₀ for this effect is 5 x 10³ IU/mg.

Endotoxin

<0.1 EU per 1 ug of the protein by the LAL method.

Amino Acid Sequence

A DNA sequence encoding Human FGF-2 Protein(#P09038) (Ala144-Ser288) was expressed.

Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Usage

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Human recombinant FGF-2 (Fibroblast Growth Factors 2) protein, AF, Tag free

For Research Use Only. Not for use in diagnostic procedures.