BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Mouse recombinant TRAIL (TNF-related apoptosis-inducing ligand) protein, AF

Catalog Number: PROTP50592-2

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Mouse recombinant TRAIL (TNF-related apoptosis-inducing ligand) protein, AF
Description	Tumor Necrosis Factor associated Apoptosis Inducing Ligand (TRAIL) is a type II membrane protein of the tumor necrosis factor (TNF) family members, moreover expressed in many adult tissues including the thymus, prostate, colon, ovary and lung. TRAIL is a 19 kDa protein containing 281 residues. Mouse TRAIL shares 70% amino acids sequence identity with human, bovine, and porcine. TRAIL to induce apoptosis in human breast carcinoma cells (MCF7) and human epithelioid carcinoma (HeLa) cell lines, by activate two death receptors of DR4 and DR5 or two decoy receptors DcR1 and DcR2.
Size	5ug,20ug,100ug
Tag	His Tag (C-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Purity

>98% as determined by SDS-PAGE.

Molecular weight

The protein has a calculated MW of 21.00 kDa. The protein migrates as 17-25 kDa under reducing condition (SDS-PAGE analysis).

Activity

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Measure by its ability to induce cytotoxicity in L929 cells in the presence of actinomycin D. The ED_{50} for this effect is <1 ng/mL.

Endotoxin

<0.1 EU per 1 ug of the protein by the LAL method.

Amino Acid Sequence

MPRGGRPQKVAAHITGITRRSNSALIPISKDGKTLGQKIESWESSRKGHSFLNHVLFRNGELVIEQEGLYYIYSQTYFRFQEAEDASKMVSKDKVRTKQLV QYIYKYTSYPDPIVLMKSARNSCWSRDAEYGLYSIYQGGLFELKKNDRIFVSVTNEHLMDLDQEASFFGAFLIN with polyhistidine tag at the C-terminus.

Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Usage

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Mouse recombinant TRAIL (TNF-related apoptosis-inducing ligand) protein, AF

For Research Use Only. Not for use in diagnostic procedures.